

# Phase segregation dynamics for the Blume-Capel model with Kac interaction.

R. Marra <sup>1</sup> and M. Mourragui <sup>2</sup>

**Abstract:** We consider the Glauber and Kawasaki dynamics for the Blume-Capel spin model with weak long range interaction on the infinite lattice: a ferromagnetic  $d$ -dimensional lattice system with the spin variable  $\sigma$  taking values in  $\{-1, 0, 1\}$  and pair Kac potential  $\gamma^d J(\gamma|i-j|)$ ,  $\gamma > 0, i, j \in \mathbb{Z}^d$ . The Kawasaki dynamics conserves the empirical averages of  $\sigma$  and  $\sigma^2$  corresponding to local magnetization and local concentration. We study the behaviour of the system under the Kawasaki dynamics on the spatial scale  $\gamma^{-1}$  and time scale  $\gamma^{-2}$ . We prove that the empirical averages converge in the limit  $\gamma \rightarrow 0$  to the solutions of two coupled equations, which are in the form of the flux gradient for the energy functional. In the case of the Glauber dynamics we still scale the space as  $\gamma^{-1}$  but look at finite time and prove in the limit of vanishing  $\gamma$  the law of large number for the empirical fields. The limiting fields are solutions of two coupled nonlocal equations. Finally, we consider a non gradient dynamics which conserves only the magnetization and get a hydrodynamic equation for it in the diffusive limit which is again in the form of the flux gradient for a suitable energy functional.

Key Words: Interacting particle and spin systems; Kac potential; hydrodynamic limits; phase segregation.

## 1. Introduction.

We consider particle models which are dynamical versions of lattice gases with Kac potentials. The Kac potentials are functions  $J_\gamma(r), r \in \mathbb{R}^d, \gamma > 0$ , such that  $J_\gamma(r) = \gamma^d J(\gamma r)$ , where  $J$  is a smooth function of compact support. They have been introduced to describe particle (or spin) systems with weak long range interaction between two particles ([KUE]). In the limit  $\gamma \rightarrow 0$  the van der Waals theory of phase transition holds exactly for these models [LP]. Here we propose to consider a Blume-Capel model with Kac interaction that we call Kac-Blume-Capel (KBC) model. The Blume-Capel model is a spin system on the lattice with nearest-neighbour interactions such that the spin variable can assume three values:  $-1, 0, 1$ . It has been introduced originally to study the  $\text{He}^3\text{-He}^4$  phase transition ([B], [C]). The structure of the phase diagram at low temperature for this model is well understood in terms of the Pirogov-Sinai theory ([BS]).

---

<sup>1</sup> Dipartimento di Fisica e Unit INFM, Universit di Roma Tor Vergata, Via della Ricerca Scientifica, 00133 Roma, Italy

<sup>2</sup> UPRESA 6085, Universit de Rouen, 76821 Mont Saint Aignan, France

The KBC model is defined by the formal Hamiltonian

$$H_\gamma(\sigma) = \frac{1}{2} \sum_{i,j \in \mathbb{Z}^d} J_\gamma(i-j)(\sigma(i) - \sigma(j))^2 - h_1 \sum_{i \in \mathbb{Z}^d} \sigma(i) - h_2 \sum_{i \in \mathbb{Z}^d} \sigma^2(i), \quad (1.1)$$

where  $h_1$  and  $h_2$  are two real parameters. In Section 2 we provide for this model the analogous of the Lebowitz-Penrose theorem, showing that in the limit  $\gamma \rightarrow 0$  the mean field theory of the Blume-Capel model ([BEG]) becomes exact. The equilibrium properties and the phase diagram of the model in the limit  $\gamma \rightarrow 0$  are very interesting. There are two order parameters characterizing the equilibrium Gibbs measure: the magnetization  $m$ , the mean value of the spin, and the concentration  $\phi$ , the mean value of the square of the spin. For inverse temperature  $\beta$  not larger than a critical value  $\beta_c$  there is a unique Gibbs measure which is indeed a Bernoulli measure (as usual for these mean field theories), while for temperatures sufficiently small (and suitable values of the parameters  $h_1$  and  $h_2$ ) the Gibbs measure is a superposition of Bernoulli measures corresponding to different values of the couple  $m, \phi$ . In particular, there is a point in the phase diagram where there are three extremal equilibrium measures, corresponding to positive, zero and negative magnetization.

We study two Markov processes in the infinite volume spin configuration space  $\Omega$  generated by self-adjoint operators in  $L^2(\Omega, \mu)$ , where  $\mu$  is a Gibbs measure for some  $\beta, h_1, h_2$  and finite  $\gamma$ : the so-called Glauber and Kawasaki dynamics. They can be described in words as follows: in the Glauber dynamics each spin at random times flips to a new value or stays unchanged with probabilities depending on the difference of energy before and after the flip. In the Kawasaki dynamics two neighboring spins at random times exchange their values, or stay unchanged, with jump rates again depending on the energy difference. The latter stochastic evolution conserves the difference and the sum between the number of spins plus and minus (respectively total magnetization and total concentration), while the former does not. Moreover, the jump rates depend on the magnetic fields  $h_1, h_2$  in the Glauber dynamics and do not in the Kawasaki one. As a consequence, all the Gibbs measures whatever are  $h_1, h_2$  are invariant for the Kawasaki dynamics, while for the Glauber dynamics the only invariant measures are the Gibbs measures with the values of  $h_1, h_2$  equal to those appearing in the jump rates.

We scale the lattice spacing by  $\gamma$  and look first at the behaviour of the system under the Glauber dynamics in the limit  $\gamma \rightarrow 0$ . We show that the empirical averages of magnetization and concentration converges weakly in probability to the solution of the set of two coupled non local equations (3.7) and (3.8) (in Section 3).

To get a definite limit in the case of the Kawasaki dynamics we have to scale also the time as  $\gamma^{-2}$  [GLP]. This is a process with two conservation laws. We prove also in this case a law of large numbers for the empirical averages of  $\sigma$  and  $\sigma^2$ , respectively magnetization and concentration. Their limits satisfy the set of two coupled non local second order integro differential equations (3.5) in Section 3. These equations can be put in a nicely form as a

gradient flow of the free energy functional  $\mathcal{F}$

$$\int dr f^0(\underline{u}(r)) + \frac{1}{2} \int dr \int dr' J(r-r') [m(r) - m(r')]^2 \quad (1.2)$$

where  $\underline{u} := (m, \phi)$  and  $f^0(\underline{u})$  is

$$f^0(\underline{u}) := -m^2 + \phi + \beta^{-1} \left[ \frac{1}{2} (m + \phi) \log(m + \phi) + \frac{1}{2} (\phi - m) \log(\phi - m) + (1 - \phi) \log(1 - \phi) - \phi \log 2 \right]. \quad (1.3)$$

The equations (3.5) become

$$\partial_t u_\alpha = \sum_{i=1}^d \sum_{\beta=1,2} \partial_i [M_{\alpha,\beta} \partial_i \frac{\delta \mathcal{F}}{\delta u_\beta}] \quad (1.4)$$

and in vectorial form

$$\partial_t \underline{u} = \nabla \cdot \left( M \nabla \frac{\delta \mathcal{F}}{\delta \underline{u}} \right), \quad (1.5)$$

where  $\frac{\delta \mathcal{F}}{\delta u_\alpha}$  denote the functional derivative of  $\mathcal{F}$  with respect to  $u_\alpha$  and  $M$  is the  $2 \times 2$  mobility matrix

$$M = \beta(1 - \phi) \begin{pmatrix} \phi + \frac{\phi^2 - m^2}{1 - \phi} & m \\ m & \phi \end{pmatrix}. \quad (1.6)$$

It is easy to see that  $\mathcal{F}$  is a Liapunov functional for (1.5). In fact

$$\frac{d}{dt} \mathcal{F} = - \sum_{i=1}^d \sum_{\alpha,\beta=1}^2 \int dr \partial_i \frac{\delta \mathcal{F}}{\delta u_\alpha} M_{\alpha,\beta} \partial_i \frac{\delta \mathcal{F}}{\delta u_\beta}. \quad (1.7)$$

The homogeneous minimizers of the functional  $\mathcal{F}$  coincide with the minimizers of  $f^0$ , which has a unique minimizer but is not convex for  $\beta$  large enough. The convex envelope of  $f^0$  is the free energy  $f$  of the KBC model at  $\gamma = 0$  and has some flat parts which single out a region  $F$  (*forbidden region*) in  $D = \{(m, \phi) : \phi \in [0, 1], m \leq \phi\}$ , the domain of definition of  $f^0$ , such that for no value of the chemical potentials  $h_1, h_2$  there is an extremal state with magnetization and concentration in  $F$ . Any Gibbs measure with averages  $m$  and  $\phi$  in  $F$  has to be a linear superposition of the extremal states with  $(m, \phi) \notin F$ .

These properties of the energy functional should allow to relate concepts of stable, unstable and metastable phases with the behaviour of the solutions of (1.5)

Glauber and Kawasaki dynamics for the Ising model with Kac potential have been investigated thoroughly ([GLP] and references therein), providing a microscopic description

of segregation phenomena. We refer for Glauber to the series [DOPT] and for Kawasaki to the papers by [GL], where the authors study the Kawasaki dynamics, with one conservation law, for the Ising model with Kac potential on a torus. Moreover they prove the hydrodynamic limit by using Radon-Nicodym derivative methods and discuss the interface motion and the segregation behaviour (see for recent developments [CCO]).

We would notice that the mean field free energy functional for the Ising model is also not convex for  $\beta$  large and, since the first order phase transition in the Ising model occurs at zero magnetic field, has a symmetric double well structure. In the KBC model instead, the phase transition (in the sense of coexistence of phases) takes place at non zero  $h_1$  and/or  $h_2$ .

Finally, we have also studied a different kind of dynamics, which is in a way intermediate between Glauber and Kawasaki in the fact that it conserves only one quantity, the magnetization. Under this dynamics a bond  $(i, j)$  (namely a couple of neighboring sites  $i$  and  $j$ ) changes its configuration  $(\sigma(i), \sigma(j))$  or stays unchanged, with probability depending on the energy difference, to a new configuration  $(\sigma'(i), \sigma'(j))$  in such a way that in each site of the bond the spin variable changes by 1 and  $\sigma(i) + \sigma(j) = \sigma'(i) + \sigma'(j)$ . Hence the magnetization stays constant during the evolution, while the number of 0's can change. For example, changes from a bond configuration  $(-1, 1)$  to a configuration  $(0, 0)$  or from  $(0, 0)$  to  $(-1, 1)$  are possible: a sort of annihilation and creation process. The jump rates are chosen to satisfy the detailed balance with respect to the Gibbs measures for the Hamiltonian (1.1) with  $h_2 = 1$ . We derive under the diffusive scaling an equation for  $m$ , while  $\phi$  on such a long time scale has already relaxed to the equilibrium and its effect can be seen in the mobility appearing in the equation for  $m$ . This dynamics is of the so called non gradient type [Sp] and the proof of the hydrodynamic limit relies on the non gradient method [V].

In Section 2 we describe the equilibrium properties of the KBC model and prove the limit  $\gamma \rightarrow 0$  for the infinite volume free energy and pressure. In Section 3 we introduce the Glauber and Kawasaki dynamics and state the main theorems, whose proofs are contained in Sections 4 and 5. In Section 6 we prove the hydrodynamic limit for the non gradient dynamics.

The proof of the hydrodynamic limit for the Kawasaki dynamics is based on the method of [GPV]. This method has been extended to infinite volume by Fritz [F], by using a bound uniform in the volume for the entropy production. In the paper by Yau [Y] a different proof of the uniform entropy production bound has been given for Ginzburg-Landau models and in [LM] this approach has been used to prove hydrodynamic limit for a class of zero range models. We follow the latter approach and prove a uniform bound for the entropy production, which is the time derivative of the entropy. Here we consider not the entropy but the relative entropy of the density of the process with respect to the Bernoulli measure  $\nu_{h_1, h_2}$  parametrized by the chemical potentials, which is not invariant for the process. Nevertheless, the bound of this production of entropy will be enough for the GPV method

to work. In fact, it is easy to show that the Kawasaki dynamics (thought of as a lattice gas dynamics) is a weak perturbation (and reduces at  $\beta = 0$  to) of the following generalized symmetric exclusion process (GSEP): each particle on the lattice jumps at random times to a nearest neighbour site  $x$  if and only if there is at most one particle in  $x$ . Hence the state of the system on times  $\gamma^{-2}$  will be very close to the invariant measures for the GSEP process, which are the Bernoulli measures  $\nu_{h_1, h_2}$ . Therefore the uniform bound for this entropy production will be sufficient to prove the hydrodynamic limit. The proof in the case of the Glauber dynamics is simpler: martingales methods are enough. In both cases it has been necessary to prove uniqueness theorems for the weak solutions of the limiting equations.

The non gradient dynamics studied in Section 6 when formulated in the language of lattice gases is a weak perturbation of a non gradient generalized simple exclusion process introduced in [KLO]. The diffusion coefficient for this process is not a constant, like in the symmetric exclusion process considered before, but a function of the density as a consequence of the non gradient character of the dynamics. We work in this case in a torus, the extension to infinite volume being more involved because of the non gradient nature of the problem. The proof is based again on the method of [GPV] and on the non gradient techniques of [V], that have to be adapted to deal with the perturbation. Also in this case, we will use as reference measure the Bernoulli measure, parametrized this time only by the magnetic field, which is not invariant for the dynamics. As a consequence, since the dynamics is non gradient, in the limiting equation there is a new term related to the solution of the non gradient problem for the unperturbed process. The presence of this term is crucial to recognize that the limiting equation is in the form of the gradient flux for a free energy functional. That is a general fact for non gradient dynamics weakly perturbed by a Kac potential, (see [GLM]). The limiting equation is

$$\partial_t m = \nabla \cdot \left( \Sigma \nabla \frac{\delta \mathcal{G}}{\delta m} \right)$$

with the energy functional  $\mathcal{G}(m(r))$  of the form

$$\int dr g^0(m(r)) + \frac{1}{2} \int dr \int dr' J(r - r') [m(r) - m(r')]^2 \quad (1.8)$$

where

$$g^0(m) := -m^2 \beta^{-1} \left[ \frac{1}{2} (m + \phi(m)) \log(m + \phi(m)) + \frac{1}{2} (\phi(m) - m) \log(\phi(m) - m) + (1 - \phi(m)) \log(1 - \phi(m)) - \phi(m) \log 2 \right] \quad (1.9)$$

and  $\phi(m) = \langle \sigma^2 \rangle_{\nu_{h_1, 0}}$  with  $h_1$  determined as a function of  $m$  via  $m(h_1) = \langle \sigma \rangle_{\nu_{h_1, 0}}$ . The mobility is given by the Einstein relation  $\Sigma = D(m)\chi(m)$ , with  $\chi$  the susceptibility

and  $D$  the diffusion coefficient, which is given by the Green-Kubo formula [KLO]. Note that  $g^0(m)$  coincide with the functional  $f^0$  in (1.3), associated to the Hamiltonian (1.1) for  $h_2 = 1$ , when evaluated in  $(m, \phi(m))$ . This is due to the fact that  $\phi$  is a fast variable under this dynamics and in the diffusive limit it relaxes to its equilibrium value  $\langle \sigma^2 \rangle_{\nu_{h_1, 0}}$ .

The convergence result that we get in this case is weaker than the one for Glauber and Kawasaki, because we are not able to prove the uniqueness of the hydrodynamic equation, due to the fact that the only regularity property known for the diffusion coefficient is continuity. Could the Lipschitz continuity for  $D$  be proven we would get the stronger convergence result.

## 2. The Kac-Blume-Capel model.

The Blume-Capel model is a model of spins with values  $0, \pm 1$  with nearest neighbours interaction, originally introduced to study the Helium phase transition. Here we introduce a model that we call Kac-Blume-Capel (KBC) model which is a model of spins taking values in  $\{-1, 0, 1\}$  on a  $d$ -dimensional lattice  $\mathbb{Z}^d$  and interacting by means of a Kac potential.

A Kac potential is a function  $J_\gamma(r)$ ,  $\gamma > 0$ , such that

$$J_\gamma(r) = \gamma^d J(\gamma r), \quad \text{for all } r \in \mathbb{R}^d,$$

where  $J \in \mathcal{C}^2(\mathbb{R}^d)$  is a non negative function supported in the unit ball, with  $\int_{\mathbb{R}^d} J(r) = 1$  and  $J(r) = J(-r)$  for all  $r \in \mathbb{R}^d$ .

The spin variable in the site  $i \in \mathbb{Z}^d$  is denoted by  $\sigma(i)$  and the infinite volume phase space is  $\{-1, 0, 1\}^{\mathbb{Z}^d}$ . A configuration is a function  $\sigma : \mathbb{Z}^d \rightarrow \{-1, 0, 1\}$ , that is an element of  $\Omega = \{-1, 0, 1\}^{\mathbb{Z}^d}$ . For any  $\Lambda \subset \mathbb{Z}^d$ , denote by  $\sigma_\Lambda$ , the restriction to  $\Lambda$  of the configuration  $\sigma$ ,  $\sigma_\Lambda = \{\sigma(i), i \in \Lambda\}$ .

The Gibbs measure, with potential  $J_\gamma(r)$  and chemical potentials  $h_1, h_2$  at inverse temperature  $\beta > 0$ , in a finite volume  $\Lambda$  and boundary condition  $\xi$  is the probability measure  $\mu_{\gamma, \Lambda}^{\beta, \xi}$  on  $\Omega$

$$\mu_{\gamma, \Lambda}^{\beta, \xi}(\sigma) = \frac{1}{Z_{\gamma, \Lambda}^{\beta, \xi}} \exp\left(-\beta H_\gamma(\sigma_\Lambda | \xi)\right),$$

where  $Z_{\gamma, \Lambda}^{\beta, \xi}$  is the normalization constant and  $H_\gamma(\sigma_\Lambda)$  is the formal Hamiltonian in a finite subset  $\Lambda$  of  $\mathbb{Z}^d$ , for the configuration  $\sigma_\Lambda$

$$H_\gamma(\sigma_\Lambda) = \frac{1}{2} \sum_{\substack{i, j \in \Lambda \\ i \neq j}} J_\gamma(i - j) (\sigma(i) - \sigma(j))^2 - h_1 \sum_{i \in \Lambda} \sigma(i) - h_2 \sum_{i \in \Lambda} \sigma^2(i), \quad (2.1)$$

The infinite volume Gibbs measure  $\mu_{\gamma, \beta}$  is a probability measure on  $\Omega$  that can be constructed by some suitable limiting procedure.

The characteristics of models with Kac potentials is that the range of the interaction is  $\gamma^{-1}$  and the strength is  $\gamma^d$ , while the total interaction with all the other spins stays finite independently of  $\gamma$ . Hence Kac potential interactions are useful to study the so-called mean field limit  $\gamma \rightarrow 0$ . The infinite volume free-energy for the Kac Ising model has been computed in the limit  $\gamma \rightarrow 0$  by Lebowitz and Penrose ([LP]), and the result agrees (and give rigorous support to) with the Van der Waals theory. The analogous result for the KBC model is

**Theorem 2.1** (*Lebowitz-Penrose limit*). *Let  $p_\gamma(\beta, h_1, h_2)$  be the pressure in the thermodynamic limit at  $\gamma > 0$ . Then*

$$\lim_{\gamma \rightarrow 0} p_\gamma(\beta, h_1, h_2) = \sup_{(m, \phi: |m| \leq \phi, \phi \leq 1)} \left[ m^2 - \phi + \beta^{-1} s(m, \phi) + h_1 m + h_2 \phi \right], \quad (2.2)$$

where  $s(m, \phi)$  is the entropy of a Bernoulli process in  $\Omega$  with average spin equal to  $m$  and average square spin equal to  $\phi$  ( $m$  is the magnetization and  $\phi$  is called concentration).

$$s(m, \phi) = -\frac{1}{2}(m + \phi) \log(m + \phi) - \frac{1}{2}(\phi - m) \log(\phi - m) - (1 - \phi) \log(1 - \phi) + \phi \log 2 \quad (2.3)$$

The free energy  $f(\beta, m, \phi)$  is defined as the Legendre transform of the pressure as

$$f(\beta, m, \phi) = \sup_{(h_1, h_2)} [h_1 m + h_2 \phi - p(\beta, h_1, h_2)] = CE \left[ -m^2 + \phi - \beta^{-1} s(m, \phi) \right], \quad (2.4)$$

where CE denotes convex envelope. The complementary result in the canonical ensemble is

**Theorem 2.2** *Define the free energy  $f_\gamma(\beta, m, \phi)$  at  $\gamma > 0$  as follows:*

*Consider the partition function in the canonical ensemble*

$$Z_\Lambda(N^-, N^0) = \sum_{\sigma \in \Gamma} \exp \left( -\beta H_\gamma(\sigma) \right), \quad (2.5)$$

where  $\Gamma$  is the set of configurations  $\{\sigma\} \in \Omega_\Lambda$  such that the number of spins  $\sigma = 1$  is fixed to be  $N^-$  and the number of spins  $\sigma = 0$  is  $N^0$ . Let  $N$  be the total number of spins in a finite volume  $\Lambda$  and put

$$\begin{aligned} m &= \frac{1}{N} \sum_{i=1}^N \sigma(i) = \frac{1}{N} [N - 2N^- N^0], \\ \phi &= \frac{1}{N} \sum_{i=1}^N (\sigma(i))^2 = \frac{1}{N} [N - N^0]. \end{aligned} \quad (2.6)$$

The free energy at  $\gamma > 0$  in the thermodynamic limit is defined as

$$\beta f_\gamma(\beta, m, \phi) := \lim_{\Lambda, N^-, N^0 \rightarrow \infty} N^{-1} \log Z_\Lambda(N^-, N^0),$$

where the limit is taken in such a way that (2.6) hold.

Then

$$\lim_{\gamma \rightarrow 0} f_\gamma(\beta, m, \phi) = CE \left[ -m^2 + \phi - \beta^{-1} s(m, \phi) \right]. \quad (2.7)$$

The proof of this theorem is similar to the one of the Lebowitz-Penrose theorem [LP], (see also [DP]) and will not be given explicitly here. We only remark that the proof of [LP] is based on a block spin renormalization procedure and the main point in the proof is writing the renormalized Hamiltonian for the block spins (whose expression will depend on the form of the interaction). Since the interaction term in the KBC model is a two body interaction like in the Ising model this part of the proof goes through in almost the same way. Obviously, the entropy will depend on which are the values of the spin and is in fact different from the one computed in [LP].

The phase structure of the model at  $\gamma = 0$  is very rich. To discuss the phase transition we can for example examine the function  $p^0 := m^2 - \phi + \beta^{-1} s(m, \phi) + h_1 m + h_2 \phi$  determining the pressure in (2.2). The extremals of the function  $p^0$  are determined by the equations

$$\begin{aligned} m &= \phi \tanh(2\beta m + \beta h_1), \\ \phi &= \exp\{-\beta(h_2 - 1)\} (1 - \phi) \cosh(2\beta m + \beta h_1). \end{aligned} \quad (2.8)$$

These equations can be solved numerically. For  $\beta \leq \bar{\beta}_c = \frac{1}{2}$  there is only one solution, while for  $\beta > \bar{\beta}_c$  the equations admit more than one solution and the function can have more than one maximum for suitable values of  $h_1$  and  $h_2$ . For  $\beta$  in the interval  $[\frac{1}{2}, \frac{3}{2}]$  there is in the plane  $\beta, h_2$  a line of second order phase transition, which changes to first order (made of triple points) at  $\beta = \bar{\beta}_c$  and  $h_2 = 1 + \frac{4}{3} \ln 2$ . The point  $\beta = \frac{3}{2}, h_1 = 0, h_2 = 1 + \frac{4}{3} \ln 2$  is called tricritical point. We refer for details to the paper [BEG]. In Fig. 1 there is the graph of  $-p^0$  as a function of  $m$  (by means of (2.8)) at a three-phases coexistence point.






Fig.1  $-p^0(m, \phi(m))$  at  $\beta = 3.76, h_1 = 0, h_2 = 0.06$ .

In Fig.2 it is shown the phase diagram in the plane  $h_1, h_2$ , for  $\beta$  large. There are three lines of phase coexistence stemming from a triple point which separate the one phase regions. In the semiplane  $h_1 > 0$  ( $h_1 < 0$ ) there is a line of coexistence of phases with positive (negative) and zero magnetization and along the line  $h_1 = 0, h_2 > k$  there is coexistence of phases with positive and negative magnetization.




Fig.2 Phase diagram at  $\beta \gg \beta_c$ .

Finally we note that the function

$$f^0 := -m^2 + \phi - \beta^{-1}s(m, \phi)$$

is not convex for  $\beta > \frac{1}{2}$ . In fact

$$Hess(f^0) = \frac{2\beta(m^2 - \phi) + 1}{(\beta^2(1 - \phi)(\phi^2 - m^2))}. \quad (2.9)$$

The Hessian is negative in the region  $\phi - m^2 > (2\beta)^{-1}$ . Hence  $f$ , the free energy at  $\gamma = 0$  defined in (2.4) as the convex envelope of  $f^0$ , has some flat parts for  $\beta > \frac{1}{2}$ , corresponding to regions in  $D = \{m, \phi : \phi \in [0, 1], m \leq \phi\}$  such that the vectorial function  $\underline{h}(m, \phi)$ ,  $\underline{h} := (h_1, h_2)$  cannot be inverted. We call this region in  $D$  forbidden region and denote it by  $F$ . A point  $(\bar{m}, \bar{\phi}) \in F$  has the property that the equations  $m(h_1, h_2) = \bar{m}, \phi(h_1, h_2) = \bar{\phi}$  cannot be solved for  $(h_1, h_2)$ .




Fig.3  $f^0(m, \phi)$  at  $\beta = 2$

Remark.

This model can also be looked at as a lattice gas of two species of particles such that in each site of the lattice there is at most one particle for each species. A way of realizing the correspondence is for example the following. Call  $\eta_b(i) = 1, 0$ ,  $\eta_r(i) = 1, 0$  the occupation number in the site  $i$  of the particles of color blue and red respectively. Then the relation  $\sigma(i) = \eta_b(i) - \eta_r(i)$  determine a lattice gas of particles blue and red with repulsive interaction between particles of the same color and attractive interaction between the same colors. Under this correspondence a configuration of particles  $\eta$  with two particles in a site  $i$  is identical to a configuration  $\zeta$  with no particles in  $i$ .

Finally we note that the relation  $\sigma(i) = \eta(i) - 1$  links the model to a lattice gas with one species of particles with at most two particles per site.

### 3. Glauber and Kawasaki dynamics.

We consider two kinds of dynamics for the spin system introduced in the previous Section: the Glauber and Kawasaki dynamics, the latter conserving both magnetization and concentration.

For  $(i, j) \in \mathbb{Z}^d \times \mathbb{Z}^d$ ,  $k \in \mathbb{Z}^d$ ,  $\sigma \in \Omega$  and any cylinder function  $F : \Omega \rightarrow \mathbb{R}$ , define

$(\nabla_{i,j}F)(\sigma)$ ,  $(\nabla_k^+F)(\sigma)$  and  $(\nabla_k^-F)(\sigma)$  by

$$\begin{aligned}(\nabla_{i,j}F)(\sigma) &= F(\sigma^{i,j}) - F(\sigma) , \\ (\nabla_k^\pm F)(\sigma) &= F(\sigma^{\pm,k}) - F(\sigma) ,\end{aligned}$$

where  $\sigma^{i,j}$  is the configuration obtained from  $\sigma$  by interchanging the value at  $i$  and  $j$  :

$$(\sigma^{i,j})(l) = \begin{cases} \sigma(l) & \text{if } l \neq i, j \\ \sigma(j) & \text{if } l = i \\ \sigma(i) & \text{if } l = j , \end{cases}$$

and  $\sigma^{\pm,k}$  is defined as

$$(\sigma^{\pm,k})(l) = \begin{cases} \sigma(l) & \text{if } l \neq k \\ \sigma(k) \pm 1 \pmod{3} & \text{if } l = k . \end{cases}$$

The Kawasaki dynamics with parameter  $\beta \geq 0$  is the unique Markov process on  $\Omega$ , whose pregenerator  $\mathbb{L}_\gamma^{K,\beta}$  acts on the cylinder functions as

$$(\mathbb{L}_\gamma^{K,\beta}f)(\sigma) = \sum_{\substack{i,j \in \mathbb{Z}^d \\ |i-j|=1}} C_\gamma^{K,\beta}(i,j;\sigma) [(\nabla_{i,j}f)(\sigma)] .$$

Here and in the following  $|\cdot|$  stands for the max norm of  $\mathbb{R}^d$ . For  $(i,j) \in \mathbb{Z}^d \times \mathbb{Z}^d$  and  $\sigma \in \Omega$ , the rate  $C_\gamma^{K,\beta}(i,j;\sigma)$  is given by

$$C_\gamma^{K,\beta}(i,j;\sigma) = \Phi\{\beta(\nabla_{i,j}H_\gamma(\sigma))\} .$$

Here  $\Phi : \mathbb{R} \rightarrow \mathbb{R}_+$  is a continuously differentiable function in a neighborhood of 0, such that  $\Phi(0) = 1$  and satisfies detailed balance condition (cf. [GL] and [GLP])

$$\Phi(E) = \exp(-E)\Phi(-E) . \tag{3.1}$$

The generator of the Glauber evolution is given by

$$(\mathbb{L}_\gamma^{G,\beta}f)(\sigma) = \sum_{i \in \mathbb{Z}^d} C_\gamma^{G,\pm}(i;\sigma) [(\nabla_i^\pm f)(\sigma)] .$$

where the rates  $C_\gamma^{G,\pm}(i;\sigma)$  are defined as

$$C_\gamma^{G,\pm}(i;\sigma) = \frac{1}{2} \frac{1}{1 + \exp\left(\beta(\nabla_i^\pm H_\gamma)\right)} .$$

corresponding to the choice  $\Phi(E) = \frac{1}{2}[1 + \exp E]^{-1}$ .

Notice that the quantities  $(\nabla_{i,j} H_\gamma)$  and  $(\nabla_i^\pm H_\gamma)$  are well defined since they involve only a finite number of non zero differences. For the proof of the existence and uniqueness of these Markov processes, we refer to Liggett [Li].

In the case of Kawasaki dynamics, if  $\beta = 0$ , the evolution reduces to a simple known process. In the setting of the lattice gas with one species of particles this dynamics is a generalized simple exclusion process GSEP [KL] with rate one, and we shall denote its pregenerator simply by  $\mathbb{L}^0$ . It differs from the usual SEP for the exclusion rule involved: in each point are allowed at most two particles. We shall see in Section 3 that the dynamics with  $\beta > 0$  is a weak perturbation of this simple exclusion. As explained in the remark in Section 3, the Kawasaki dynamics can also be interpreted as the motion of two species of particles, moving as a symmetric simple exclusion process with rate one, with the exclusion rule  $\eta_b + \eta_r \leq 1$ , such that also jumps exchanging colors between neighbour sites are allowed. If such jumps are forbidden the system becomes a non gradient system and the diffusion coefficient in this case is different from one ([Q]).

Since the Kawasaki dynamics conserves magnetization and concentration the invariant measures will be Gibbs measures parametrized by two chemical potentials. It is useful to introduce the invariant measures for the exclusion process GSEP, which are Bernoulli measures depending on two parameters. For each positive integer  $n$ , denote by  $\Lambda_n \subset \mathbb{Z}^d$  the sublattice of size  $(2n+1)^d$ ,  $\Lambda_n = \{-n, \dots, n\}^d$ . For  $A = (a, b) \in [-1, 1] \times [0, 1]$ , we define  $\bar{\nu}_A$  as the product measure on  $\Omega$  with chemical potential  $A$  such that, for all positive integer  $n$ , the restriction  $\bar{\nu}_{A,n}$  of  $\bar{\nu}_A$  to  $\Omega_n$  is given by

$$d\bar{\nu}_{A,n} = Z_{A,n}^{-1} \exp \left\{ a \sum_{i \in \Lambda_n} \sigma(i) + b \sum_{i \in \Lambda_n} \sigma^2(i) \right\},$$

where  $Z_{A,n}$  is the normalization constant. For  $(a, b) \in [-1, 1] \times [0, 1]$  let  $m = m(a, b)$  (resp.  $\phi = \phi(a, b)$ ) be the expectation of  $\sigma(0)$  (resp.  $\sigma^2(0)$ ) under  $\bar{\nu}_{A,n}$ :

$$\begin{cases} m(a, b) = \mathbf{E}^{\bar{\nu}_{A,n}}(\sigma(0)) \\ \phi(a, b) = \mathbf{E}^{\bar{\nu}_{A,n}}(\sigma^2(0)). \end{cases}$$

Observe that the function  $\Psi$  defined on  $] - 1, 1[ \times ] 0, 1[$  by  $\Psi(a, b) = (m, \phi)$  is a bijection from  $] - 1, 1[ \times ] 0, 1[$  to  $I = \left\{ (m, \phi) : 0 < \phi < 1, -1 < m < \phi \right\}$ . For every  $P = (m, \phi) \in I$ , we denote by  $\nu_{P,n}$  the product measure such that

$$\begin{cases} m = \mathbf{E}^{\nu_{P,n}}[\sigma(0)] \\ \phi = \mathbf{E}^{\nu_{P,n}}[\sigma^2(0)]. \end{cases} \quad (3.2)$$

We take  $\gamma^{-1}$ , the range of the interaction, as macroscopic space unit and consider the limit  $\gamma \rightarrow 0$ . We want to establish for both Kawasaki and Glauber dynamics a law of large numbers for the empirical fields corresponding to magnetization and concentration.

In the Glauber case we look at the behaviour of the fields for finite time, while in the Kawasaki case the relevant time scale is  $\gamma^{-2}$ . Fix a sequence of probability measures  $(\mu^\gamma)_\gamma$ , associated to the same initial profile  $(m_0, \phi_0) : \mathbb{R}^d \times \mathbb{R}^d \longrightarrow I$  in the following sense

$$\lim_{\gamma \rightarrow 0} \mu^\gamma \left\{ \left[ \left| \gamma^d \sum_{i \in \mathbb{Z}^d} U(i\gamma) \sigma(i) - \int_{\mathbb{R}^d} U(x) m_0(x) dx \right| + \left| \gamma^d \sum_{i \in \mathbb{Z}^d} V(i\gamma) \sigma(i)^2 - \int_{\mathbb{R}^d} V(x) \phi_0(x) dx \right| \right] > \delta \right\} = 0, \quad (3.3)$$

for every continuous functions  $U, V : \mathbb{R}^d \longrightarrow \mathbb{R}$  with compact support, and every  $\delta > 0$ .

Denote by  $\mathbf{P}_{\mu^\gamma}^{(K),\beta}$  (resp.  $\mathbf{P}_{\mu^\gamma}^{(G),\beta}$ ) the probability measure on the path space  $D(\mathbb{R}_+, \Omega)$  corresponding to the Markov process  $(\sigma(t, \cdot))_{t \geq 0}$  with the generator  $\gamma^{-2} \mathbb{L}_\gamma^{K,\beta}$  (resp.  $\mathbb{L}_\gamma^{G,\beta}$ ), and starting from  $\mu^\gamma$ , and by  $\mathbf{E}_{\mu^\gamma}^{(K),\beta}$  (resp.  $\mathbf{E}_{\mu^\gamma}^{(G),\beta}$ ) the expectation with respect to  $\mathbf{P}_{\mu^\gamma}^{(K),\beta}$  (resp.  $\mathbf{P}_{\mu^\gamma}^{(G),\beta}$ ). Denote by  $\mathcal{C}_K^2(\mathbb{R}^d)$  the space of real twice continuously differentiable functions with compact support.

The main result of this Section is the following theorem.

**Theorem 3.1** *Under (3.3), for any  $\delta > 0$ ,  $t \geq 0$  and  $U, V \in \mathcal{C}_K^2(\mathbb{R}^d)$  the following holds*

(i) *Kawasaki dynamics.*

$$\lim_{\gamma \rightarrow 0} \mathbf{P}_{\mu^\gamma}^{(K),\beta} \left\{ \left[ \left| \gamma^d \sum_{i \in \mathbb{Z}^d} U(i\gamma) \sigma(t, i) - \int_{\mathbb{R}^d} U(x) m(t, x) dx \right| + \left| \gamma^d \sum_{i \in \mathbb{Z}^d} V(i\gamma) \sigma(t, i)^2 - \int_{\mathbb{R}^d} V(x) \phi(t, x) dx \right| \right] > \delta \right\} = 0, \quad (3.4)$$

where  $(m, \phi)$  is the unique weak solution of

$$\begin{aligned} \partial_t m &= \nabla \cdot \left[ \nabla m - 2\beta(\phi - (m)^2)(\nabla J * m) \right], \\ \partial_t \phi &= \nabla \cdot \left[ \nabla \phi - 2\beta m(1 - \phi)(\nabla J * m) \right], \\ m(0, \cdot) &= m_0, \quad \phi(0, \cdot) = \phi_0 \end{aligned} \quad (3.5)$$

where  $*$  denotes the convolution on the spatial variable.

(ii) *Glauber dynamics.*

$$\lim_{\gamma \rightarrow 0} \mathbf{P}_{\mu^\gamma}^{(G), \beta} \left\{ \left[ \left| \gamma^d \sum_{i \in \mathbb{Z}^d} U(i\gamma) \sigma(t, i) - \int_{\mathbb{R}^d} U(x) m^{(G)}(t, x) dx \right| \right. \right. \\ \left. \left. + \left| \gamma^d \sum_{i \in \mathbb{Z}^d} V(i\gamma) \sigma(t, i)^2 - \int_{\mathbb{R}^d} V(x) \phi^{(G)}(t, x) dx \right| \right] > \delta \right\} = 0, \quad (3.6)$$

where  $(m^{(G)}, \phi^{(G)})$  is the unique weak solution of

$$\begin{cases} \partial_t \begin{pmatrix} m^{(G)} \\ \phi^{(G)} \end{pmatrix} = \mathcal{G}^\beta \begin{pmatrix} m^{(G)} \\ \phi^{(G)} \end{pmatrix} := \begin{pmatrix} \mathcal{G}_1^\beta(m^{(G)}, \phi^{(G)}) \\ \mathcal{G}_2^\beta(m^{(G)}, \phi^{(G)}) \end{pmatrix} \\ \begin{pmatrix} m^{(G)}(0, \cdot) \\ \phi^{(G)}(0, \cdot) \end{pmatrix} = \begin{pmatrix} m_0 \\ \phi_0 \end{pmatrix}. \end{cases} \quad (3.7)$$

and  $\mathcal{G}_1^\beta, \mathcal{G}_2^\beta$  are defined as

$$\begin{aligned} \mathcal{G}_1^\beta(m, \phi) &= \frac{1}{4} \left\{ \tanh \frac{\beta}{2}(\alpha + h') + \tanh \frac{\beta}{2}(\alpha - h') \right\} \\ &\quad + \frac{1}{4} \frac{m}{2} \left\{ \tanh \frac{\beta}{2}(\alpha + h') - \tanh \frac{\beta}{2}(\alpha - h') \right\} - \frac{3}{4} m \\ &\quad + \frac{\phi}{4} \left\{ 2 \tanh \beta \alpha - \frac{1}{2} \tanh \frac{\beta}{2}(\alpha + h') - \frac{1}{2} \tanh \frac{\beta}{2}(\alpha - h') \right\} \\ \mathcal{G}_2^\beta(m, \phi) &= \frac{1}{4} \left\{ \tanh \frac{\beta}{2}(\alpha + h') - \tanh \frac{\beta}{2}(\alpha - h') \right\} \\ &\quad + \frac{1}{4} \frac{m}{2} \left\{ \tanh \frac{\beta}{2}(\alpha - h') + \tanh \frac{\beta}{2}(\alpha + h') \right\} \\ &\quad - \frac{1}{4} \frac{\phi}{2} \left\{ \tanh \frac{\beta}{2}(\alpha + h') - \tanh \frac{\beta}{2}(\alpha - h') \right\} + \frac{1}{4} (2 - 3\phi). \end{aligned} \quad (3.8)$$

Here

$$\alpha = 2m * J + h_1, \quad h' = h_2 - \int_{\mathbb{R}^d} J(x) dx = h_2 - 1.$$

and  $*$  denotes the convolution.

The limiting equations for the Kawasaki dynamics can be rewritten in a nice form as a gradient flux associated with the local mean field free energy functional. Put  $\underline{u} := (m, \phi)$ . Define the free energy functional as

$$\mathcal{F}(\underline{u}) := -\frac{1}{\beta} \int dr s(\underline{u}(r)) - \int dr \int dr' J(r - r') m(r) m(r')$$

and the mobility matrix as

$$M = -\beta[Hess(s)]^{-1} = \beta(1 - \phi) \begin{pmatrix} \phi + \frac{\phi^2 - m^2}{1 - \phi} & m \\ m & \phi \end{pmatrix}.$$

Then equations (3.5) become

$$\partial_t \underline{u} = \nabla \left( M \nabla \frac{\delta \mathcal{F}}{\delta \underline{u}} \right)$$

Writing the free energy functional  $\mathcal{F}$  in the equivalent form

$$\int dr f^0(\underline{u}(r)) + \frac{1}{2} \int dr \int dr' J(r - r') [m(r) - m(r')]^2$$

with

$$f^0(\underline{u}) := -m^2 + \phi - \beta^{-1} s(\underline{u})$$

we see that  $\mathcal{F}$  reduces for homogeneous profiles of magnetization and concentration to the non convex free energy  $f^0$  of the KBC model, so that the stationary homogeneous solutions of the equations (3.5) coincide with the solutions of (2.8). Moreover,  $\mathcal{F}$  is a Liapunov functional for the evolution, namely is decreasing in time along the solutions of the equations (3.5). This follows from (1.7) and the positivity of the matrix  $M$ .

On the contrary, in the Glauber case the limiting equations (3.7)-(3.8) are rather messy. It is not even known if the energy functional is a Liapunov functional: we have only numerical evidence.

The region  $D$  in the plane  $(m, \phi)$ , such that  $D = \{m, \phi : 0 \leq \phi \leq 1, |m| \leq \phi, \}$  can be partitioned for any fixed  $\beta \geq \bar{\beta}_c$  in three parts:

- a) the unstable region  $U = \{(m, \phi) \in F : \phi - m^2 \geq (2\beta)^{-1}\}$ , where  $F$  is the forbidden region defined after (2.9).
- b) the metastable region  $M = \{(m, \phi) \in F : \phi - m^2 < (2\beta)^{-1}\}$
- c) the stable region  $D - (U \cup M)$ .

The segregation phenomena may appear by choosing an initial datum corresponding to total magnetization and concentration in the unstable region. One expects that a stationary solution of the equations (3.5) with this initial condition be unstable.

#### 4. Dirichlet form estimates for Kawasaki dynamics.

The proof of Theorem 3.1 is based on a priori estimates uniform in the volume for the entropy and the Dirichlet form, which are given in this section. For each positive integer  $n$  and a measure  $\mu$  on  $\Omega_n = \{-1, 0, 1\}^{\Lambda_n}$ , we denote by  $\mu_n$  the marginal of  $\mu$  on  $\Omega_n$ ,

$$\mu_n(\xi) = \mu\{\sigma : \sigma(i) = \xi(i) \text{ for } |i| \leq n\} \text{ for each } \xi \in \Omega_n.$$

For a chemical potential  $P$ , and a positive integer  $n$ , we denote by  $s_n(\mu_n|\nu_{P,n})$  the relative entropy of  $\mu_n$  with respect to  $\nu_{P,n}$

$$s_n(\mu_n|\nu_{P,n}) = \sup_{U \in C_b(\Omega_n)} \left\{ \int U(\sigma) d\mu_n(\sigma) - \log \int e^{U(\sigma)} d\nu_{P,n}(\sigma) \right\}.$$

In this formula  $C_b(\Omega_n)$  stands for the space of all functions on  $\Omega_n$ . Since the measure  $\nu_{P,n}$  gives a positive probability to each configuration, all the measures on  $\Omega_n$  are absolutely continuous with respect to  $\nu_{P,n}$  and we have an explicit formula for the entropy :

$$s_n(\mu_n|\nu_{P,n}) = \int \log(f_n(\sigma)) d\mu_n(\sigma),$$

where  $f_n$  is the probability density of  $\mu_n$  with respect to  $\nu_{P,n}$ . Notice that by the entropy convexity and since  $\sup_{\sigma} \sup_i |\sigma(i)|$  is finite, we have

$$s_n(\mu_n|\nu_{P,n}) \leq C_0 n^d \tag{4.1}$$

for some constant  $C_0$  that depends on  $P$  (cf. [KL]).

Define the Dirichlet form  $D_n(\mu_n|\nu_{P,n})$  of the measure  $\mu_n$  with respect to  $\nu_{P,n}$  associated to the exclusion process by

$$\begin{aligned} D_n(\mu_n|\nu_{P,n}) &= - \int \sqrt{f_n}(\sigma) (\mathbb{L}_n^0 \sqrt{f_n})(\sigma) d\nu_{P,n}(\sigma) \\ &= \sum_{\substack{i,j \in \Lambda_n \\ |i-j|=1}} I_{i,j}(f_n), \end{aligned}$$

where  $I_{i,j}(\cdot)$  is given by

$$I_{i,j}(f_n) = - \int \sqrt{f_n}(\sigma) (\mathbb{L}_{i,j}^0 \sqrt{f_n})(\sigma) d\nu_{P,n}(\sigma)$$

and  $\mathbb{L}_n^0$  is the restriction of the process to the box  $\Lambda_n$

$$\mathbb{L}_n^0 = \sum_{\substack{i,j \in \Lambda_n \\ |i-j|=1}} \mathbb{L}_{i,j}^0.$$

Here for a bond  $(i,j) \in \mathbb{Z}^d \times \mathbb{Z}^d$ ,  $\mathbb{L}_{i,j}^0$  stands for the piece of generator associated to the exchange of spins between sites  $i$  and  $j$  for the exclusion process.

Define the entropy  $\mathcal{S}(\mu|\nu_P)$  and the Dirichlet form  $\mathcal{D}(\mu|\nu_P)$  of a measure  $\mu$  on  $\Omega$  with respect to  $\nu_P$  as

$$\begin{aligned} \mathcal{S}(\mu|\nu_P) &= \gamma \sum_{n \geq 1} s_n(\mu_n|\nu_{P,n}) e^{-\gamma n}, \\ \mathcal{D}(\mu|\nu_P) &= \gamma \sum_{n \geq 1} D_n(\mu_n|\nu_{P,n}) e^{-\gamma n}. \end{aligned}$$



Notice that by (4.1), there exists a positive constant  $C$  depending on  $P$  such that for any probability measure  $\mu$  on  $\Omega$

$$\mathcal{S}(\mu|\nu_P) \leq C\gamma^{-d}. \quad (4.2)$$

Through this section we consider Kawasaki dynamics with fixed parameter  $\beta > 0$  and with fixed scaling parameter  $\gamma^{-1}$ . We shall denote by  $(S_\gamma^{K,\beta}(t))_{t \geq 0}$  the semigroup associated to the generator  $\gamma^{-2}\mathbb{L}_\gamma^{K,\beta}$  (that is the semigroup of Kawasaki dynamics with parameter  $\beta$ , accelerated by  $\gamma^{-2}$ ). For a measure  $\mu$  on  $\Omega$  we shall denote by  $\mu^{K,\beta}(t)$  the time evolution of the measure  $\mu$  under the semigroup  $S_\gamma^{K,\beta}$ :  $\mu^{K,\beta}(t) = \mu S_\gamma^{K,\beta}(t)$ .

When  $\beta = 0$ , the process reduces to the generalized simple exclusion process (Lemma 4.1), and in particular the product measures are invariant for the generator  $\mathbb{L}_\gamma^{K,0}$ . In this case, by using the methods of [F] and [Y] one can get entropy and the Dirichlet form estimates uniform in the volume, for the entropy of processes evolving in large finite volumes and then extend them by lower semi-continuity to the infinite system. For  $\beta \neq 0$  product measures are no more invariant for the generator, but it is possible to take advantage from the fact that the process is a weak perturbation of the exclusion one (Lemma 4.1 below) and adapt the Fritz's approach to that case without considering an approximation of the infinite volume dynamics. Notice that from (4.2) there is no need for an initial condition on the entropy in Theorem 3.1.

**Lemma 4.1** *For any  $i \in \mathbb{Z}^d$ , unit vector  $e \in \mathbb{Z}^d$  and  $\sigma \in \Omega$*

$$\begin{aligned} C_\gamma^{K,\beta}(i, i+e; \sigma) &= 1 - \gamma\beta[\sigma(i+e) - \sigma(i)]\gamma^d \sum_{\ell \in \mathbb{Z}^d} (e \cdot \nabla J)(\gamma(i-\ell))\sigma(\ell) \\ &\quad - \gamma^{d+1}\beta[\sigma(i+e) - \sigma(i)]^2(e \cdot \nabla J)(0) + O(\gamma^2) \\ &= 1 + \beta O(\gamma). \end{aligned}$$

*Proof.* By definition of  $H_\gamma$ , for all  $i, j \in \mathbb{Z}^d$  and  $\sigma \in \Omega$

$$\begin{aligned} (\nabla_{i,j} H_\gamma)(\sigma) &= 2[\sigma(i) - \sigma(j)]\gamma^d \sum_{\ell \in \mathbb{Z}^d} [J_\gamma(i-\ell) - J_\gamma(j-\ell)]\sigma(\ell) \\ &\quad + 2[\sigma(i) - \sigma(j)]^2\gamma^d [J_\gamma(i-j) - J_\gamma(0)]. \end{aligned} \quad (4.3)$$

To prove the lemma, it is enough to remark that the conditions imposed on  $\Phi$  imply that  $\Phi'(0) = -\frac{1}{2}$  (cf. [GL]) and to use Taylor expansion.  $\square$

We get the following estimate for the Dirichlet form in the infinite volume:

**Theorem 4.2** *There exists a positive finite constant  $C_1$  that depends on  $P$ ,  $t$  and  $\beta$  such that*

$$\int_0^t \mathcal{D}(\mu^{K,\beta}(\tau)|\nu_P) d\tau \leq C_1 \gamma^{2-d}.$$

The strategy of the proof is to introduce suitable entropy and Dirichlet form in finite volume and bound the corresponding entropy production in terms of the finite volume Dirichlet form times  $\gamma^2$  uniformly in the volume (Lemma 4.2). Then, the a priori bound on the entropy (4.2) allows to get the estimate.

Fix a measure  $\mu$  on  $\Omega$  and a chemical potential  $P$ . For every  $t \geq 0$  and positive integer  $n$ , denote by  $f_n^t$  the probability density of  $(\mu^{K,\beta}(t))_n$  with respect to  $\nu_{P,n}$ . To simplify the notation, we denote respectively by  $s_n(f_n^t)$  and  $D_n(f_n^t)$  the entropy and the Dirichlet form of  $(\mu^{K,\beta}(t))_n$  with respect to  $\nu_{P,n}$ . For all positive integer  $M$ , let  $M_\gamma$  be defined by  $M_\gamma = N_\gamma^2 + M$ , where  $N_\gamma = \llbracket \gamma^{-1} \rrbracket$  stands for the integer part of  $\gamma^{-1}$ . Define respectively the entropy  $\mathcal{S}_{M_\gamma}(\cdot)$  and the Dirichlet form  $\mathcal{D}_{M_\gamma}(\cdot)$  with finite sum by

$$\begin{aligned} \mathcal{S}_{M_\gamma}(\mu^{K,\beta}(t)|\nu_P) &= \gamma \sum_{n=1}^{M_\gamma} s_n(f_n^t) e^{-n\gamma}, \\ \mathcal{D}_{M_\gamma}(\mu^{K,\beta}(t)|\nu_P) &= \gamma \sum_{n=1}^{M_\gamma} D_n(f_n^t) e^{-n\gamma}. \end{aligned}$$

**Lemma 4.3** *There exist positive and finite constants  $A_0$  and  $A_1$  that depend on  $P$  and  $\beta$  such that, for all positive  $M$*

$$\partial_t \mathcal{S}_{M_\gamma}(\mu^{K,\beta}(t)|\nu_P) \leq -\gamma^{-2} A_0 \mathcal{D}_{M_\gamma}(\mu^{K,\beta}(t)|\nu_P) + A_1 \gamma^{-d}. \quad (4.4)$$

Before proving the Lemma we conclude the proof of Theorem 4.2.

*Proof of Theorem 4.2.* Integrate (4.4) from 0 to  $t$ , let  $M \uparrow \infty$  and use (4.2).  $\square$

*Proof of Lemma 4.3.* We drop the indices in  $\mathbb{L}_\gamma^{K,\beta}$  and denote the generator simply by  $\mathbb{L}$ , For all positive integer  $k$  denote by  $\mathbb{L}_k$  the restriction of the generator  $\mathbb{L}$  to the box  $\Lambda_k$ . For a subset  $A \subset \Lambda$  of  $\mathbb{Z}^d$ , and a function  $h$  in  $L^1(\nu_{P,\Lambda})$ , let  $\langle h \rangle_A$  be the function on  $\{-1, 0, 1\}^{\Lambda \setminus A}$  obtained by integrating  $h$  over the coordinates  $\{\sigma(x) : x \in A\}$  with respect to  $\nu_{P,\Lambda}$ . When  $A = \Lambda_{n+m+1} - \Lambda_n$ , we shall denote this expectation simply by  $\langle h \rangle_{\Lambda_n^m}$ .

With this notation, we can verify that  $f_n^t$  satisfy the equation

$$\partial_t f_n^t = \gamma^{-2} \left\langle \mathbb{L}_{n+1}^* f_{n+N_\gamma+1}^t \right\rangle_{\Lambda_n^{n+N_\gamma}}, \quad (4.5)$$

where for a positive integer  $k$ ,  $\mathbb{L}_k^*$  represents the adjoint operator of  $\mathbb{L}_k$  in  $L^2(\nu_{P,\Lambda_k})$ . By relation (4.5) and the explicit formula for the entropy we have that

$$\begin{aligned}\partial_t s_n(f_n^t) &= \gamma^{-2} \int f_{n+N_\gamma+1}^t \mathbb{L}_n \log(f_n^t) d\nu_{P,n+N_\gamma+1} \\ &\quad + \gamma^{-2} \int f_{n+N_\gamma+1}^t (\partial L_{n+1}) \log(f_n^t) d\nu_{P,n+N_\gamma+1} \\ &:= \Omega_n^1 + \Omega_n^2.\end{aligned}\tag{4.6}$$

The first term  $\Omega_n^1$  in the right hand side of the last inequality corresponds to the exchanges in the interior of  $\Lambda_n$ , while the second term  $\Omega_n^2$  is associated to exchanges at the boundary

$$(\partial L_{n+1})(f) = \sum_{\substack{i \in \Lambda_n, j \notin \Lambda_n \\ |i-j|=1}} C_\gamma^{K,\beta}(i,j;\sigma) \left[ (\nabla_{i,j} f)(\sigma) \right].$$

The proof is divided in three steps. In the first two steps we estimate  $\Omega_n^1$  and  $\Omega_n^2$  and in the third one we prove (4.4) .

*First step (bound of  $\Omega_n^1$ ).*

Fix a bond  $(i,j) \in \Lambda_n \times \Lambda_n$  such that  $|i-j|=1$ , denote by  $\mathbb{L}_{i,j}$  the one bond generator corresponding to the exchange of spins between  $i$  and  $j$  and let  $F_n^{i,j}(\sigma)$  be the function defined by  $F_n^{i,j}(\sigma) = \left\langle \frac{C_\gamma^{K,\beta}(i,j;\cdot)}{f_n^t(\sigma)} f_{n+N_\gamma+1}^t(\cdot) \right\rangle_{\Lambda_n^{n+N_\gamma}}$ . We have

$$\gamma^{-2} \int f_{n+N_\gamma+1}^t \mathbb{L}_{i,j} \log(f_n^t) d\nu_P = \gamma^{-2} \int F_n^{i,j}(\sigma) f_n^t(\sigma) \log \left\{ \frac{f_n^t(\sigma^{i,j})}{f_n^t(\sigma)} \right\} d\nu_P(\sigma).$$

Using the basic inequality

$$a(\log b - \log a) \leq -(\sqrt{a} - \sqrt{b})^2 + (b - a)\tag{4.7}$$

for positive  $a$  and  $b$ , the right hand side of the last expression is bounded by

$$-\gamma^{-2} \int F_n^{i,j}(\sigma) \left[ \sqrt{f_n^t(\sigma^{i,j})} - \sqrt{f_n^t(\sigma)} \right]^2 d\nu_P + \gamma^{-2} \int F_n^{i,j}(\sigma) \left[ f_n^t(\sigma^{i,j}) - f_n^t(\sigma) \right] d\nu_P.\tag{4.8}$$

Observe that for all function  $h$  and positive integers  $n$  and  $m$ ,  $\langle h_{n+m+1} \rangle_{\Lambda_n^{n+m}} = h_n$ . In particular, using Lemma 4.1 we have that

$$\left| F_n^{i,j}(\sigma) - 1 \right| \leq B\gamma.$$

With this remark, and since the measure  $\nu_P$  is invariant for the exclusion process, (4.8) is bounded above by

$$-\gamma^{-2}(1 - B\gamma) \int \left[ \sqrt{f_n^t(\sigma^{i,j})} - \sqrt{f_n^t(\sigma)} \right]^2 d\nu_P + B\gamma^{-1} \int \left| f_n^t(\sigma^{i,j}) - f_n^t(\sigma) \right| d\nu_P.$$

Using the elementary inequality  $2ab \leq A^{-1}a^2 + Ab^2$ , the second term of the last inequality is bounded by

$$\frac{A}{2}\gamma^{-2}I_{i,j}(f_n^t) + \frac{B}{2A} \int \left[ \sqrt{f_n^t(\sigma^{i,j})} + \sqrt{f_n^t(\sigma)} \right]^2 d\nu_P \leq \frac{A}{2}\gamma^{-2}I_{i,j}(f_n^t) + 2\frac{B^2}{A},$$

where we used in the last inequality, Schwartz inequality and the fact that  $f_n^t$  is the probability density with respect to  $\nu_P$ . Choosing  $A$  small enough, and taking the sum over all  $i, j \in \Lambda_n$  such that  $|i - j| = 1$ , we get

$$\Omega_n^1 \leq -C_0 D_n(f_n^t) + C'_0 n^d \quad (4.9)$$

for some positive constants  $C_0$  and  $C'_0$ .

*Second step (bound of  $\Omega_n^2$ ).*

Fix a bond  $(i, j) \in \Lambda_n \times \Lambda_n^c$ , such that  $|i - j| = 1$  and decompose  $\mathbb{I}_{i,j}$  in three terms,

$$\mathbb{I}_{i,j} = \mathbb{I}_{i,j}^{(0,1)} + \mathbb{I}_{i,j}^{(-1,0)} + \mathbb{I}_{i,j}^{(-1,1)}, \quad (4.10)$$

where for  $(l, m) \in \{(0, 1), (-1, 0), (-1, 1)\}$ ,  $\mathbb{I}_{i,j}^{(l,m)}$  is given by

$$\begin{aligned} (\mathbb{I}_{i,j}^{(l,m)} g)(\sigma) &= r_{i,j}^{(l,m)}(\sigma) C_\gamma^{K,\beta}(i, j; \sigma) \left[ g(\mathbf{T}_{m-l}^{j,i} \sigma) - g(\sigma) \right] \\ &\quad + r_{j,i}^{(l,m)}(\sigma) C_\gamma^{K,\beta}(i, j; \sigma) \left[ g(\mathbf{T}_{m-l}^{i,j} \sigma) - g(\sigma) \right]. \end{aligned}$$

Here for  $\iota = 1, 2$ ,  $(i, j) \in \mathbb{Z}^d \times \mathbb{Z}^d$  and a configuration  $\sigma$ ,  $\mathbf{T}_\iota^{i,j} \sigma$  is defined by  $\mathbf{T}_\iota^{i,j} \sigma = \sigma - \iota \delta_i + \iota \delta_j$ , and  $r_{i,j}^{(l,m)}(\sigma) = 1_{\{\sigma(i)=l, \sigma(j)=m\}}$ . For  $k \in \mathbb{Z}^d$ ,  $\delta_k$  is the configuration with spin 1 at site  $k$  and none elsewhere, and addition of two configurations is defined coordinate by coordinate.

The term  $\Omega_n^2$  in (4.6) can be written as a sum of terms  $\Omega_{i,j}^2$  associated to the bond  $(i, j)$ . The decomposition (4.10) induces an analogous decomposition for  $\Omega_{i,j}^2$ . We study explicitly only the one corresponding to  $\mathbb{I}_{i,j}^{(-1,1)}$ , that we denote by  $\Omega_{i,j}^{(-1,1)}$ . The other two terms are dealt with in the same way.

$$\Omega_{i,j}^{(-1,1)} = \gamma^{-2} \int f_{n+N_\gamma+1}^t(\sigma) \mathbb{I}_{i,j}^{(-1,1)} \log(f_n^t(\sigma)) d\nu_P.$$

Let  $F_1^{i,j}$  and  $F_2^{i,j}$  be defined by

$$\begin{aligned} F_1^{i,j}(\sigma) &= 1_{\{\sigma(j)=1\}} C_\gamma^{K,\beta}(i, j; \sigma) f_{n+N_\gamma+1}^t(\sigma), \\ F_2^{i,j}(\sigma) &= 1_{\{\sigma(j)=1\}} C_\gamma^{K,\beta}(i, j; \sigma^{j,i}) f_{n+N_\gamma+1}^t(\sigma^{j,i}). \end{aligned}$$

By changing variables,  $\Omega_{i,j}^{(-1,1)}$  can be rewritten as

$$\gamma^{-2} \int 1_{\{\sigma(i)=-1\}} \left\{ \langle F_1^{i,j}(\sigma) \rangle_{\Lambda_n^{n+N_\gamma}} - \langle F_2^{i,j}(\sigma) \rangle_{\Lambda_n^{n+N_\gamma}} \right\} \log \left\{ \frac{f_n^t(\sigma + 2\delta_i)}{f_n^t(\sigma)} \right\} d\nu_P(\sigma). \quad (4.11)$$

Since we have that  $(a-b)(\log c - \log d)$  is negative for  $a, b, c$  and  $d$  positive real numbers, unless  $a \geq b$  and  $c \geq d$  or  $a \leq b$  and  $c \leq d$ , we may introduce in the last integral the indicator function of the set  $E_n^1 \cup E_n^2$ , where

$$E_n^1 = \left\{ \sigma : \langle F_1^{i,j}(\sigma) \rangle_{\Lambda_n^{n+N_\gamma}} \geq \langle F_2^{i,j}(\sigma) \rangle_{\Lambda_n^{n+N_\gamma}}, f_n^t(\sigma + 2\delta_i) \geq f_n^t(\sigma) \right\},$$

$$E_n^2 = \left\{ \sigma : \langle F_1^{i,j}(\sigma) \rangle_{\Lambda_n^{n+N_\gamma}} \leq \langle F_2^{i,j}(\sigma) \rangle_{\Lambda_n^{n+N_\gamma}}, f_n^t(\sigma + 2\delta_i) \leq f_n^t(\sigma) \right\}.$$

We shall consider separately the integral on  $E_n^1$  and  $E_n^2$ , and we call  $\Omega_4$  (resp.  $\Omega_5$ ) the integral on the set  $E_n^1$  (resp.  $E_n^2$ ). We consider first the integral on  $E_n^1$  and rewrite it as the sum of two other terms

$$\begin{aligned} \Omega_4 &= \gamma^{-2} \int 1_{\{\sigma(i)=-1\}} \left\{ \langle F_1^{i,j}(\sigma) \rangle_{\Lambda_n^{n+N_\gamma}} - \langle F_3^{i,j}(\sigma) \rangle_{\Lambda_n^{n+N_\gamma}} \right\} \log \left\{ \frac{f_n^t(\sigma + 2\delta_i)}{f_n^t(\sigma)} \right\} 1_{E_n^1} d\nu_P(\sigma) \\ &\quad + \gamma^{-2} \int 1_{\{\sigma(i)=-1\}} \left\{ \langle F_3^{i,j}(\sigma) \rangle_{\Lambda_n^{n+N_\gamma}} - \langle F_2^{i,j}(\sigma) \rangle_{\Lambda_n^{n+N_\gamma}} \right\} \log \left\{ \frac{f_n^t(\sigma + 2\delta_i)}{f_n^t(\sigma)} \right\} 1_{E_n^1} d\nu_P(\sigma). \end{aligned}$$

where  $F_3^{i,j}$  is defined by

$$F_3^{i,j}(\sigma) = 1_{\{\sigma(j)=1\}} C_\gamma^{K,\beta}(i, j; \sigma^{j,i}) f_{n+N_\gamma+1}^t(\sigma).$$

Applying Lemma 4.1 below, we obtain that the first line of the last expression is of order  $\gamma^{-1}$ . Indeed, observe that we have for all configurations  $\sigma$

$$\begin{aligned} &\gamma^{-2} \langle F_1^{i,j}(\sigma) \rangle_{\Lambda_n^{n+N_\gamma}} - \langle F_3^{i,j}(\sigma) \rangle_{\Lambda_n^{n+N_\gamma}} \\ &= \gamma^{-2} \langle 1_{\{\sigma(j)=1\}} [C_\gamma^{K,\beta}(i, j; \sigma) - C_\gamma^{K,\beta}(i, j; \sigma^{i,j})] f_{n+N_\gamma+1}^t(\sigma) \rangle_{\Lambda_n^{n+N_\gamma}} \\ &\leq C_2 \gamma^{-1} \langle f_{n+N_\gamma+1}^t(\sigma) \rangle_{\Lambda_n^{n+N_\gamma}} = C_2 \gamma^{-1} f_n^t(\sigma) \end{aligned}$$

for some positive constant  $C_2$ , and on the set  $E_n^1$ , we have that  $f_n^t(\sigma + 2\delta_i) \geq f_n^t(\sigma)$ . Hence, from inequality (4.7), the change of variables and the fact that  $f_n^t$  is a probability density with respect to  $\nu_P$ , the first term of  $\Omega_4$  is bounded by

$$C_2 \gamma^{-1} \int 1_{\{\sigma(i)=-1\}} |f_n^t(\sigma + 2\delta_i) - f_n^t(\sigma)| d\nu_P(\sigma) \leq C'_2 \gamma^{-1}$$

for some positive constant  $C'_2$  that depends on  $P$  and  $\beta$ . We estimate now the second term of  $\Omega_4$ . Since on  $E_n^1$  we have  $f_n^t(\sigma + 2\delta_i) \geq f_n^t(\sigma)$ , we may replace the indicator function on  $E_n^1$  by the indicator function on the set  $E_n^3$  defined by

$$E_n^3 = \left\{ \sigma : \langle F_3^{i,j}(\sigma) \rangle_{\Lambda_n^{n+N_\gamma}} \geq \langle F_2^{i,j}(\sigma) \rangle_{\Lambda_n^{n+N_\gamma}}, f_n^t(\sigma + 2\delta_i) \geq f_n^t(\sigma) \right\}.$$

Since for all positive  $x$ ,  $\log x \leq 2(\sqrt{x} - 1)$ , and since on the set  $E_n^3$ , we have that  $\langle F_3^{i,j}(\sigma) \rangle_{\Lambda_n^{n+N_\gamma}} \geq \langle F_2^{i,j}(\sigma) \rangle_{\Lambda_n^{n+N_\gamma}}$ , the second integral of  $\Omega_4$  with indicator function on  $E_n^3$  is less than or equal to

$$2\gamma^{-2} \int 1_{\{\sigma(i)=-1\}} \left\{ \langle F_3^{i,j}(\sigma) \rangle_{\Lambda_n^{n+N_\gamma}} - \langle F_2^{i,j}(\sigma) \rangle_{\Lambda_n^{n+N_\gamma}} \right\} \left\{ \frac{\sqrt{f_n^t(\sigma + 2\delta_i)}}{\sqrt{f_n^t(\sigma)}} - 1 \right\} 1_{E_n^3} d\nu_P(\sigma).$$

By the elementary inequality  $2xy \leq \frac{\gamma^{-1}}{\alpha} x^2 + \frac{\alpha}{\gamma^{-1}} y^2$  and since  $(A-B) = (\sqrt{A} - \sqrt{B})(\sqrt{A} + \sqrt{B})$  we have that for every positive  $A, B, a, b$  and for every positive  $\alpha$ ,

$$2(A-B)((b/a) - 1) \leq \frac{\gamma^{-1}}{\alpha} (\sqrt{A} - \sqrt{B})^2 + \frac{\alpha}{\gamma^{-1}} (\sqrt{A} + \sqrt{B})^2 ((b/a) - 1)^2.$$

In particular the last integral is bounded above by

$$\begin{aligned} & \frac{\gamma^{-3}}{\alpha} \int 1_{\{\sigma(i)=-1\}} \left\{ \sqrt{\langle F_3^{i,j}(\sigma) \rangle_{\Lambda_n^{n+N_\gamma}}} - \sqrt{\langle F_2^{i,j}(\sigma) \rangle_{\Lambda_n^{n+N_\gamma}}} \right\}^2 d\nu_P(\sigma) \\ & + \alpha \gamma^{-1} \int 1_{\{\sigma(i)=-1\}} \left\{ \sqrt{\langle F_3^{i,j}(\sigma) \rangle_{\Lambda_n^{n+N_\gamma}}} + \sqrt{\langle F_2^{i,j}(\sigma) \rangle_{\Lambda_n^{n+N_\gamma}}} \right\}^2 \\ & \quad \times \left\{ \frac{\sqrt{f_n^t(\sigma + 2\delta_i)}}{\sqrt{f_n^t(\sigma)}} - 1 \right\}^2 1_{E_n^3} d\nu_P(\sigma). \end{aligned} \quad (4.12)$$

The first line of this expression is bounded above by the one bond Dirichlet form. It is equal to

$$\begin{aligned} & \frac{\gamma^{-3}}{N_\gamma \alpha} \sum_{m=n+N_\gamma+1}^{n+2N_\gamma} \int \left\{ \sqrt{\langle r_{i,j}^{(-1,1)}(\sigma) C_\gamma^{K,\beta}(i,j; \sigma^{i,j}) f_m^t(\sigma^{j,i}) \rangle_{\Lambda_n^m}} \right. \\ & \quad \left. - \sqrt{\langle r_{i,j}^{(-1,1)}(\sigma) C_\gamma^{K,\beta}(i,j; \sigma^{i,j}) f_m^t(\sigma) \rangle_{\Lambda_n^m}} \right\}^2 d\nu_P(\sigma), \end{aligned}$$

which, by Schwartz inequality, is bounded by

$$\begin{aligned} & \frac{\gamma^{-3}}{N_\gamma \alpha} \sum_{m=n+N_\gamma+1}^{n+2N_\gamma} \int \langle r_{i,j}^{(-1,1)}(\sigma) C_\gamma^{K,\beta}(i,j; \sigma^{i,j}) \{ \sqrt{f_m^t(\sigma^{j,i})} - \sqrt{f_m^t(\sigma)} \}^2 \rangle_{\Lambda_n^m} d\nu_P(\sigma) \\ & \leq C_4 \frac{\gamma^{-2}}{\alpha} \sum_{m=n+N_\gamma+1}^{n+2N_\gamma} \int \langle r_{i,j}^{(-1,1)}(\sigma) \{ \sqrt{f_m^t(\sigma^{j,i})} - \sqrt{f_m^t(\sigma)} \}^2 \rangle_{\Lambda_n^m} d\nu_P(\sigma) \\ & = C_4 \frac{\gamma^{-2}}{\alpha} \sum_{m=n+N_\gamma+1}^{n+2N_\gamma} \int r_{i,j}^{(-1,1)}(\sigma) \{ \sqrt{f_m^t(\sigma^{j,i})} - \sqrt{f_m^t(\sigma)} \}^2 d\nu_P(\sigma), \end{aligned}$$

where we used the fact that there exists a positive constant  $C_4$ , such that  $C_\gamma^{K,\beta}(i, j; \sigma) \leq C_4$ , for all configuration  $\sigma$ .

Finally the second term of (4.12) is bounded by

$$\begin{aligned} & 4\gamma^{-1}\alpha \int 1_{\{\sigma(i)=-1\}} \frac{\langle F_3^{i,j}(\sigma) \rangle_{\Lambda_n^{n+N_\gamma}}}{f_n^t(\sigma)} \left\{ \sqrt{f_n^t(\sigma + 2\delta_i)} - \sqrt{f_n^t(\sigma)} \right\}^2 d\nu_P(\sigma) \\ & \leq 4C_4\gamma^{-1}\alpha \int 1_{\{\sigma(i)=-1\}} \left\{ \sqrt{f_n^t(\sigma + 2\delta_i)} - \sqrt{f_n^t(\sigma)} \right\}^2 d\nu_P(\sigma) \leq C_5\gamma^{-1}\alpha \end{aligned}$$

for some positive constant  $C_5$  that depends on  $P$ . We have changed variables and used the fact that  $f_n^t$  is a probability density with respect to  $\nu_P$ .

Collecting the above inequalities, we get the following bound for  $\Omega_4$ . For any positive  $\alpha$

$$\begin{aligned} \Omega_4 & \leq \gamma^{-1}(C'_2 + C_5\alpha) \\ & + C_4 \frac{\gamma^{-2}}{\alpha} \sum_{m=n+N_\gamma+1}^{n+2N_\gamma} \int r_{i,j}^{(-1,1)}(\sigma) \left\{ \sqrt{f_m^t(\sigma^{j,i})} - \sqrt{f_m^t(\sigma)} \right\}^2 d\nu_P(\sigma). \end{aligned} \quad (4.13)$$

The term  $\Omega_5$  will be handled in an analogous way. It can be rewritten as

$$\begin{aligned} \Omega_5 & = \gamma^{-2} \int 1_{\{\sigma(i)=-1\}} \left\{ \langle F_2^{i,j}(\sigma) \rangle_{\Lambda_n^{n+N_\gamma}} - \langle F_4^{i,j}(\sigma) \rangle_{\Lambda_n^{n+N_\gamma}} \right\} \log \left\{ \frac{f_n^t(\sigma)}{f_n^t(\sigma + 2\delta_i)} \right\} 1_{E_n^2} d\nu_P(\sigma) \\ & + \gamma^{-2} \int 1_{\{\sigma(i)=-1\}} \left\{ \langle F_4^{i,j}(\sigma) \rangle_{\Lambda_n^{n+N_\gamma}} - \langle F_1^{i,j}(\sigma) \rangle_{\Lambda_n^{n+N_\gamma}} \right\} \log \left\{ \frac{f_n^t(\sigma)}{f_n^t(\sigma + 2\delta_i)} \right\} 1_{E_n^2} d\nu_P(\sigma) \end{aligned}$$

with

$$F_4^{i,j} = 1_{\{\sigma(j)=1\}} C_\gamma^{K,\beta}(i, j; \sigma) f_{n+N_\gamma+1}^t(\sigma^{j,i}).$$

By the same arguments used to estimate  $\Omega_4$ , we obtain by exchanging the role of  $f_n^t(\sigma)$  and  $f_n^t(\sigma + 2\delta_i)$ ,

$$\begin{aligned} \Omega_5 & \leq \gamma^{-1}(C'_2 + C_5\alpha) \\ & + C_4 \frac{\gamma^{-2}}{\alpha} \sum_{m=n+N_\gamma+1}^{n+2N_\gamma} \int r_{j,i}^{(-1,1)}(\sigma) \left\{ \sqrt{f_m^t(\sigma^{i,j})} - \sqrt{f_m^t(\sigma)} \right\}^2 d\nu_P(\sigma) \end{aligned}$$

for all positive  $\alpha$ . Therefore, taking advantage of this last inequality and of (4.13), we get

$$\begin{aligned} & \gamma^{-2} \int f_{n+N_\gamma+1}^t(\sigma) \mathbb{L}_{i,j}^{(-1,1)} \log(f_n^t(\sigma)) d\nu_P \leq 2\gamma^{-1}(C'_2 + C_5\alpha) \\ & + C_4 \frac{\gamma^{-2}}{\alpha} \sum_{m=n+N_\gamma+1}^{n+2N_\gamma} \int r_{i,j}^{(-1,1)}(\sigma) \left\{ \sqrt{f_m^t(\sigma^{j,i})} - \sqrt{f_m^t(\sigma)} \right\}^2 d\nu_P(\sigma). \\ & + C_4 \frac{\gamma^{-2}}{\alpha} \sum_{m=n+N_\gamma+1}^{n+2N_\gamma} \int r_{j,i}^{(-1,1)}(\sigma) \left\{ \sqrt{f_m^t(\sigma^{i,j})} - \sqrt{f_m^t(\sigma)} \right\}^2 d\nu_P(\sigma). \end{aligned}$$

To conclude this step, we have just to sum over  $\{(0, 1), (-1, 0), (-1, 1)\}$  and over  $\{(i, j) \in \Lambda_n \times \Lambda_n^c : |i - j| = 1\}$ . We obtain

$$\Omega_n^2 \leq 6n^{d-1}\gamma^{-1}(C'_2 + C_5\alpha) + C_4 \frac{\gamma^{-2}}{\alpha} \sum_{m=n+N_\gamma}^{n+2N_\gamma} I_{i,j}(f_m^t) \quad (4.14)$$

for any positive  $\alpha$ .

*Third step (Proof of (4.4)).*

From (4.6), (4.9) and (4.14), for all positive  $n$

$$\begin{aligned} \partial_t s_n(t) &\leq -C_0 D_n(f_n^t) + C'_0 n^d + 6n^{d-1}\gamma^{-1}(C'_2 + C_5\alpha) \\ &\quad + C_4 \frac{\gamma^{-2}}{\alpha} \sum_{(i,j) \in \Lambda_n \times \Lambda_n^c} \sum_{m=n+N_\gamma+1}^{n+2N_\gamma} I_{i,j}(f_m^t). \end{aligned}$$

Multiply both sides of this inequality by  $e^{-n\gamma}$ , sum over  $1 \leq n \leq M_\gamma$  and for  $\alpha$  large enough. We get for some positive constants  $A_0$  and  $A_1$

$$\begin{aligned} \partial_t \mathcal{S}_M(t) &\leq -A_0 \mathcal{D}_n(f_n^t) + A_1 \gamma^{-d} \\ &\quad + C'_4 \gamma^{-2} \sum_{n=M_\gamma-2N_\gamma+1}^{M_\gamma} e^{-n\gamma} \sum_{m=1}^{N_\gamma} \sum_{(i,j) \in \Lambda_n \times \Lambda_n^c} I_{i,j}(f_{m+n+N_\gamma}^t). \end{aligned}$$

To conclude the proof of Lemma 4.3, it remains to observe that the third term of the right hand side of the last inequality is bounded by  $Const.\gamma^{-d}$ .  $\square$

**Corollary 4.4** *For all  $K > 0$ ,*

$$\int_0^t D_{KN_\gamma}(f_{KN_\gamma}^\tau) d\tau \leq 2C_1 e^{K+1} \gamma^{2-d}.$$

*Proof.* Fix  $K > 0$  and  $\tau \in [0, t]$ , since by Schwartz inequality  $n \mapsto D_n(f_n^\tau)$  is a nondecreasing function we have

$$\begin{aligned} D_{KN_\gamma}(f_{KN_\gamma}^\tau) &\leq \frac{1}{N_\gamma} \sum_{m=1}^{N_\gamma} D_{KN_\gamma+m}(f_{KN_\gamma+m}^\tau) \\ &\leq e^{K+1} \frac{1}{N_\gamma} \sum_{m=1}^{N_\gamma} D_{KN_\gamma+m}(f_{KN_\gamma+m}^\tau) e^{-(KN_\gamma+m)\gamma} \\ &\leq 2e^{K+1} \mathcal{D}(\mu^{K,\beta}(\tau) | \nu_P). \end{aligned}$$

$\square$



## 5. Hydrodynamic limits for Kawasaki and Glauber dynamics.

In this section we prove Theorem 3.1. Let  $\mathcal{C}_K(\mathbb{R}^d)$  denote the space of real continuous functions (with compact support) and denote by  $\mathcal{M}$  the space of signed measures on  $\mathbb{R}^d$  with total variation bounded by 1 equipped with the weak\* topology induced by  $\mathcal{C}_K(\mathbb{R}^d)$  via  $\langle \mu, U \rangle = \int U d\mu$ .

Given a configuration  $\sigma(t, \cdot)$ , we define the empirical measures  $\pi_t^{1,\gamma}(\sigma(t, \cdot)) = \pi_t^{1,\gamma}$  and  $\pi_t^{2,\gamma}(\sigma(t, \cdot)) = \pi_t^{2,\gamma}$  by

$$(\pi_t^{1,\gamma}, \pi_t^{2,\gamma}) = \left( \gamma^d \sum_{i \in \mathbb{Z}^d} \sigma(t, i) \delta_{\gamma i}, \gamma^d \sum_{i \in \mathbb{Z}^d} (\sigma(t, i))^2 \delta_{\gamma i} \right),$$

where  $\delta_{\gamma i}$  is the Dirac mass at the macroscopic site  $\gamma i$ . We shall denote in the sequel  $\sigma(t, i)$  by  $\sigma_t(i)$  and  $(\sigma(t, i))^2$  by  $\sigma(t, i)^2$ .

*Kawasaki dynamics.*

First of all, in order to prove (3.4) it is enough to show that, for any positive time  $t$ , any functions  $U, V \in \mathcal{C}_K(\mathbb{R}^d)$  and  $\delta > 0$ ,

$$\lim_{\gamma \rightarrow 0} \mathbf{P}_{\mu^\gamma}^{(K),\beta} \left\{ \left| \langle \pi_t^{1,\gamma}, U \rangle + \langle \pi_t^{2,\gamma}, V \rangle - \int_{\mathbb{R}^d} \left( m^{(K)}(t, x)U(x) + \phi^{(K)}(t, x)V(x) \right) dx \right| > \delta \right\} = 0.$$

where  $(m^{(K)}(\cdot, \cdot), \phi^{(K)}(\cdot, \cdot))$  is a weak solution of the hydrodynamic equations (3.5).

Fix a parameter  $\beta > 0$  and consider the Kawasaki dynamics at  $\beta$  positive. For a fixed time interval  $[0, T]$ , we denote by  $\mathbf{P}_{\mu^\gamma}^{(K)}$  the law of the process  $(\sigma_t)_{t \in [0, T]}$  accelerated by  $\gamma^{-2}$  on the space  $D([0, T], \Omega)$  and by  $Q_{\mu^\gamma}^{(K)}$  the law of the process  $(\pi_t^{1,\gamma}, \pi_t^{2,\gamma})_{t \in [0, T]}$  on the space  $D([0, T], \mathcal{M}^2)$  with initial distribution  $\mu^\gamma$ . The law of large numbers for the empirical measures  $\pi_t^{1,\gamma}$  and  $\pi_t^{2,\gamma}$  follows ([GPV]) from the weak convergence of the probability measures  $Q_{\mu^\gamma}^{(K)}$  to a probability  $Q^{(K)}$  concentrated on the deterministic trajectory  $(\pi^1(t, dx), \pi^2(t, dx)) = (m^{(K)}(t, x)dx, \phi^{(K)}(t, x)dx)$ , where  $(m^{(K)}(\cdot, \cdot), \phi^{(K)}(\cdot, \cdot))$  is a weak solution of the hydrodynamic equations (3.5). The proof of this result requires tightness, identification of the limit and uniqueness of the weak solution of the limiting equation.

**Lemma 5.1** *Tightness. The sequence  $(Q_{\mu^\gamma}^{(K)})_\gamma$  is a tight family and all its limit points  $Q^*$  are such that*

$$\begin{aligned} Q^* \left\{ (\pi^1, \pi^2) : (\pi^1(t, dx), \pi^2(t, dx)) &= (\pi^1(t, x)dx, \pi^2(t, x)dx) \right\} = 1, \\ Q^* \left\{ (\pi^1, \pi^2) : -1 \leq \pi^1(t, x) \leq 1, 0 \leq \pi^2(t, x) \leq 1 \right\} &= 1. \end{aligned}$$

The proof of 5.1 is very simple since  $\sup_\sigma \sup_i |\sigma(i)| < \infty$  and therefore is omitted.

**Lemma 5.2** *All limit points  $Q^*$  of the sequence  $(Q_{\mu^\gamma}^{(K)})_\gamma$  are concentrated on weak solutions of equation (3.5).*

Finally, the law of large numbers follows from the uniqueness of the weak solution of equations (3.5), whose proof is given in Lemma 5.4 below.

*Proof of Lemma 5.2.* Denote by  $\mathcal{C}_K^{1,2}([0, T] \times \mathbb{R}^d)$  the space of compact support functions  $V : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$ , twice continuously differentiable on space with continuous derivative in time. Fix a function  $U = (U^1, U^2)$  such that  $U^1(t, x) = U_t^1(x)$  and  $U^2(t, x) = U_t^2(x)$  are in  $\mathcal{C}_K^{1,2}([0, T] \times \mathbb{R}^d)$ , and consider the martingale  $M_t^U$  defined by

$$M_t^U = \sum_{n=1}^2 \left\{ \langle \pi_t^{n,\gamma}, U_t^n \rangle - \langle \pi_0^{n,\gamma}, U_0^n \rangle - \int_0^t \left( \partial_s + \gamma^{-2} \mathbb{I}_\gamma^{K,\beta} \right) \langle \pi_s^{n,\gamma}, U_s^n \rangle ds \right\}$$

with the quadratic variation  $\mathcal{N}_t^U$  given by

$$\begin{aligned} \mathcal{N}_t^U = (M_t^U)^2 - \int_0^t & \left\{ \gamma^{-2} \mathbb{I}_\gamma^{K,\beta} \left( \sum_{n=1}^2 \langle \pi_s^{n,\gamma}, U_s^n \rangle \right)^2 \right. \\ & \left. - 2 \left( \sum_{n=1}^2 \langle \pi_s^{n,\gamma}, U_s^n \rangle \right) \gamma^{-2} \mathbb{I}_\gamma^{K,\beta} \left( \sum_{n=1}^2 \langle \pi_s^{n,\gamma}, U_s^n \rangle \right) \right\} ds. \end{aligned}$$

Denote by  $\{e_1, \dots, e_d\}$  the orthonormal basis of  $\mathbb{R}^d$ , and observe that for all  $i \in \mathbb{Z}^d$ ,  $\sigma \in \Omega$  and  $k = 1, \dots, d$  we have  $C_\gamma^{K,\beta}(i + e_k, i; \sigma) = \tau_{e_k} C_\gamma^{K,\beta}(i, i - e_k; \sigma)$ , where  $\tau_{e_k}$  is the space shift by  $e_k$  acting on  $\Omega$ . Hence, a spatial summation by parts permits to rewrite the integral term of  $M_t^U$  as

$$\begin{aligned} & \sum_{n=1}^2 \int_0^t \langle \pi_s^{n,\gamma}, \partial_s U_s^n \rangle ds \\ & - \sum_{n=1}^2 \int_0^t \gamma^{d-1} \sum_{i \in \mathbb{Z}^d} \sum_{k=1}^d C_\gamma^{K,\beta}(i + e_k, i; \sigma_s) [\sigma_s(i + e_k)^n - \sigma_s(i)^n] (\partial_k^\gamma U_s^n)(\gamma i) ds. \end{aligned}$$

Here  $\partial_k^\gamma$  represents the discrete derivative in the  $k$ -th direction:

$$(\partial_k^\gamma V)(\gamma i) = \gamma^{-1} [V(\gamma(i + e_k)) - V(\gamma i)].$$

Notice that the conditions imposed on  $\Phi$  imply that  $\Phi'(0) = -1/2$  (cf. [GL]), in particular, using Lemma 4.1 and a second summation by parts, we may rewrite the second term of the last integral as

$$\begin{aligned} & \sum_{n=1}^2 \left\{ \gamma^d \sum_{i \in \mathbb{Z}^d} \sum_{k=1}^d \int_0^t ds \left\{ (\sigma_s(i)^n) \gamma^{-2} [U_s^n(\gamma(i + e_k)) + U_s^n(\gamma(i - e_k)) - 2U_s^n(\gamma i)] \right\} \right\} \\ & + \frac{\beta}{2} \sum_{n=1}^2 \left\{ \gamma^d \sum_{i \in \mathbb{Z}^d} \sum_{k=1}^d \int_0^t ds \left\{ (\partial_k^\gamma U_s^n)(\gamma i) (\tau_i g^n(\sigma)) (\pi_s^{1,\gamma} * \partial_k^\gamma J)(\gamma i) \right\} \right\} + o_\gamma(1), \end{aligned}$$

where  $o_\gamma(1)$  is a random variable that converges to 0 with  $\gamma$  and  $*$  stands for the convolution in the spatial variable. For  $n = 1, 2$ , the functions  $g^n$  are defined by

$$g^n(\sigma) = [\sigma(e_k) - \sigma(0)] [\sigma(e_k)^n - \sigma(0)^n].$$

This time, however, it is not the density fields themselves that appear in the second term of the last expression but another local function of the configuration. Following the methods of Guo-Papanicolaou-Varadhan [GPV], the main step in proving the hydrodynamic equations is to replace this local function by another function of the density fields in order to close the equations. For a cylinder function  $\Psi$ , we denote its expectation with respect to the measure  $\nu_{(m,\phi)} = \nu_P$  defined in (3.2) by  $\tilde{\Psi}(m, \phi)$ :

$$\tilde{\Psi}(m, \phi) = \int \Psi(\sigma) d\nu_P(\sigma)$$

and for a positive integer  $\ell$  and  $i \in \mathbb{Z}^d$ , denote the empirical mean densities on a box of size  $(2\ell + 1)^d$  centered at  $i$  by  $(\mathbf{A}^{1,\ell}\sigma)(i)$  and  $(\mathbf{A}^{2,\ell}\sigma)(i)$ :

$$\left( (\mathbf{A}^{1,\ell}\sigma)(i), (\mathbf{A}^{2,\ell}\sigma)(i) \right) = \left( \frac{1}{(2\ell + 1)^d} \sum_{|i-j|\leq\ell} \sigma(j), \frac{1}{(2\ell + 1)^d} \sum_{|i-j|\leq\ell} \sigma(j)^2 \right).$$

**Lemma 5.3** *For every cylinder function  $\Psi$  and every  $V : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$  with compact support*

$$\limsup_{\varepsilon \rightarrow 0} \limsup_{\gamma \rightarrow 0} \mathbf{E}_{\mu_\gamma}^{(K)} \left[ \gamma^d \sum_{i \in \mathbb{Z}^d} \int_0^T ds |V_s(\gamma i)| \left\{ \left| (2\varepsilon N_\gamma + 1)^{-d} \sum_{|j-i|\leq\varepsilon N_\gamma} \tau_j \Psi(\sigma_s) - \tilde{\Psi} \left( (\mathbf{A}^{1,\varepsilon N_\gamma}\sigma_s)(i), (\mathbf{A}^{2,\varepsilon N_\gamma}\sigma_s)(i) \right) \right| \right\} \right] = 0,$$

where  $N_\gamma$  is the integer part of  $\gamma^{-1}$ .

Since the support of the function  $V$  is compact, by corollary 4.4 the proof of this Lemma is very similar to the one usually used in finite volume. Nevertheless, we shall give a sketch of its proof at the end of this subsection.

Let us go on with the proof of Lemma 5.2. Now, by Lemma 5.3 and Taylor expansion applied to the functions  $U^1$ ,  $U^2$  and  $J$ , the integral term of the martingale  $M_t^U$  can be written as

$$\sum_{n=1}^2 \sum_{k=1}^d \int_0^t ds \left\{ \langle \pi_s^{n,\gamma}, (\partial_s U_s^n + \partial_k^2 U_s^n) \rangle + \frac{\beta}{2} \sum_{i \in \mathbb{Z}^d} \left[ (\partial_k U_s^n)(\gamma i) (\pi_s^{1,\gamma} * \partial_k J)(\gamma i) \tilde{g}^n \left( (\mathbf{A}^{1,\varepsilon N_\gamma}\sigma)(i), (\mathbf{A}^{2,\varepsilon N_\gamma}\sigma)(i) \right) \right] \right\} + o_{\gamma,\varepsilon}(1),$$

where  $\partial_k$  and  $\partial_k^2$  represent the first and the second derivatives in the  $k$ -th direction, and  $o_{\gamma,\varepsilon}(1)$  is a random variable that converges to 0 when  $\gamma \rightarrow 0$  and  $\varepsilon \rightarrow 0$ .

Moreover, remark that

$$\begin{aligned}\tilde{g}^1\left((\mathbf{A}^{1,\varepsilon N_\gamma}\sigma)(i), (\mathbf{A}^{2,\varepsilon N_\gamma}\sigma)(i)\right) &= 2\left[(\mathbf{A}^{2,\varepsilon N_\gamma}\sigma)(i) - \left((\mathbf{A}^{1,\varepsilon N_\gamma}\sigma)(i)\right)^2\right], \\ \tilde{g}^2\left((\mathbf{A}^{1,\varepsilon N_\gamma}\sigma)(i), (\mathbf{A}^{2,\varepsilon N_\gamma}\sigma)(i)\right) &= 2(\mathbf{A}^{1,\varepsilon N_\gamma}\sigma)(i)\left[1 - (\mathbf{A}^{2,\varepsilon N_\gamma}\sigma)(i)\right].\end{aligned}$$

On the other hand, a simple computation shows that the quadratic variation of the martingale  $M_t^U$  is equal to

$$\begin{aligned}\mathcal{N}_t^U = (M_t^U)^2 - \gamma^{2d} \sum_{\substack{i,j \\ |i-j|=1}} \int_0^t \left\{ C_\gamma^{K,\beta}(i,j;\sigma_s) \right. \\ \left. \times \left( \sum_{n=1}^2 \gamma^{-1} [U_s^n(\gamma i) - U_s^n(\gamma j)] [\sigma_s(i)^n - \sigma_s(j)^n] \right)^2 \right\} ds\end{aligned}$$

and then it vanishes as  $\gamma$  goes to 0. By Doob's inequality, for every  $\delta > 0$ ,

$$\lim_{\gamma \rightarrow 0} \mathbf{P}_{\mu^\gamma}^{(K)} \left\{ \sup_{0 \leq t \leq T} |M_t^U| > \delta \right\} = 0.$$

Therefore, collecting the above arguments and using Lemma 5.1, we obtain that any limit point  $Q^*$  of the sequence  $(Q_{\mu^\gamma}^{(K)})_\gamma$  is such that

$$\begin{aligned}Q^* \left\{ (m, \phi) : \left| \langle m_T, U_T \rangle - \langle m_0, U_0 \rangle - \int_0^T \langle m_s, (\partial_s U_s + \Delta U_s) \rangle ds \right. \right. \\ \left. \left. + \langle \phi_T, V_T \rangle - \langle \phi_0, V_0 \rangle - \int_0^T \langle \phi_s, (\partial_s V_s + \Delta V_s) \rangle ds \right. \right. \\ \left. \left. - \beta \sum_{k=1}^d \int_0^T \langle (m_s * (\partial_k J)) \left( (\phi_s * \alpha_\varepsilon) - (m_s * \alpha_\varepsilon)^2 \right), (\partial_k U_s) \rangle ds \right. \right. \\ \left. \left. - \beta \sum_{k=1}^d \int_0^T \langle (m_s * (\partial_k J)) (m_s * \alpha_\varepsilon) (1 - (\phi_s * \alpha_\varepsilon)), (\partial_k V_s) \rangle ds \right| > \delta \right\} = 0\end{aligned}$$

for any  $\delta > 0$  and  $U, V \in \mathcal{C}_K^{1,2}([0, T] \times \mathbb{R}^d)$ , where  $\alpha_\varepsilon$  is defined as  $\alpha_\varepsilon(x) = (2\varepsilon)^{-d} \mathbf{1}_{[-\varepsilon, \varepsilon]^d}(x)$ . Let  $\varepsilon$  go to 0 and by arbitrariness of  $\delta$  we obtain the statement of Lemma 5.2.  $\square$

*Proof of Lemma 5.3.* To simplify the notation, for  $\sigma \in \Omega$ , denote by  $\mathbf{V}^{N_\gamma, \varepsilon}(\sigma)$  the expression

$$\mathbf{V}^{N_\gamma, \varepsilon}(\sigma) = \frac{1}{(2\varepsilon N_\gamma + 1)^d} \left| \sum_{|j| \leq \varepsilon N_\gamma} \Psi(\sigma) - \tilde{\Psi} \left( (\mathbf{A}^{1,\varepsilon N_\gamma}\sigma)(0), (\mathbf{A}^{2,\varepsilon N_\gamma}\sigma)(0) \right) \right|.$$

Fix  $M > 0$  such that  $[-M, M]^d$  contains the support of  $V_s$  for all  $s \in [0, T]$ . We have

$$\mathbf{E}_{\mu^\gamma}^{(K)} \left[ \gamma^d \sum_{i \in \mathbb{Z}^d} \int_0^T |V_s(\gamma i)| \tau_i \mathbf{V}^{N_\gamma, \varepsilon}(\sigma_s) ds \right] \leq \|V\|_\infty \mathbf{E}_{\mu^\gamma}^{(K)} \left[ \gamma^d \sum_{i \in \Lambda_{MN_\gamma}} \int_0^T \tau_i \mathbf{V}^{N_\gamma, \varepsilon}(\sigma_s) ds \right].$$

Denote  $\bar{f}^T = \frac{1}{T} \int_0^T f_{(M+2)N_\gamma}^s ds$ , where  $f_{(M+2)N_\gamma}^s$  is the probability density with respect to  $\nu_{P, (M+2)N_\gamma} = \nu_P^{M, N_\gamma}$  of the restriction of the measure  $\mu^{K, \beta}(s)$  to the box  $\Lambda_{(M+2)N_\gamma}$  (see section 3). Since the function  $\sum_{i \in \Lambda_{MN_\gamma}} \tau_i \mathbf{V}^{N_\gamma, \varepsilon}(\sigma)$  depends on the configuration  $\sigma$  only through the variables  $\{\sigma(k) : k \in \Lambda_{(M+1)N_\gamma}\}$ , by Fubini's theorem, Dirichlet form convexity and Corollary 4.4, there exists a positive constant  $C$  that depends on  $M, \beta$  and  $P$  such that the right hand side of the last inequality is bounded by

$$\left\{ T \|V\|_\infty \int \gamma^d \sum_{i \in \Lambda_{MN_\gamma}} \tau_i \mathbf{V}^{N_\gamma, \varepsilon}(\sigma) \bar{f}^T(\sigma) d\nu_P^{M, N_\gamma}(\sigma) - A \gamma^{d-2} D_{(M+2)N_\gamma}(\bar{f}^T) \right\} + AC$$

for all positive  $A$ . It follows that, in order to prove Lemma 5.3 it is enough to show that for each positive  $A$

$$\limsup_{\varepsilon \rightarrow 0} \limsup_{\gamma \rightarrow 0} \sup \left\{ \int \gamma^d \sum_{i \in \Lambda_{MN_\gamma}} \tau_i \mathbf{V}^{N_\gamma, \varepsilon}(\sigma) f(\sigma) d\nu_P^{M, N_\gamma}(\sigma) - A \gamma^{d-2} D_{(M+2)N_\gamma}(f) \right\} = 0,$$

where the supremum is carried over all probability densities  $f$  with respect to  $\nu_P^{M, N_\gamma}$ .

The proof of this limit relies on the usual one and two blocks estimates (cf. [GPV] or [KOV]) and therefore is omitted.  $\square$

**Lemma 5.4** (Uniqueness). *For any  $T > 0$ , the equation (3.5) has a unique weak solution in the class  $L^\infty([0, T] \times \mathbb{R}^d) \times L^\infty([0, T] \times \mathbb{R}^d)$ .*

*Proof.* The proof follows the arguments in [GL] adapted to the infinite volume case. For a positive time  $t > 0$ ,  $f \in \mathcal{C}_K(\mathbb{R}^d)$  and  $\varepsilon > 0$ , let  $\mathcal{H}_{t, \varepsilon}^f : [0, t] \times \mathbb{R}^d \rightarrow \mathbb{R}$  be defined by

$$\mathcal{H}_{t, \varepsilon}^f(s, x) = (f * h_{t+\varepsilon-s})(x),$$

where  $h_{t+\varepsilon-s}(\cdot)$  is the heat kernel given by

$$h_{t+\varepsilon-s}(x) = (2\pi(t+\varepsilon-s))^{-d/2} \exp \left\{ -\frac{1}{4(t+\varepsilon-s)} \sum_{k=1}^d (x_k)^2 \right\}.$$

In the Appendix it is proven that  $\mathcal{H}_{t,\varepsilon}^f$  solves the equation  $\partial_t \rho = \Delta \rho$  on  $[0, t] \times \mathbb{R}^d$  and that

$$\begin{aligned} \sum_{k=1}^d \int_0^t \left| \langle \partial_k \mathcal{H}_{t,\varepsilon}^f(s, \cdot) \rangle \right| ds &\leq C_1 (\sqrt{t+\varepsilon} - \sqrt{\varepsilon}) \|f\|_1 \\ &\leq C_1 \sqrt{t} \|f\|_1, \end{aligned} \quad (5.1)$$

where  $C_1$  is a positive constant that depends on  $d$  and  $\partial_k$  is the first derivative in the  $k$ -th direction.

Let us consider  $(m, \phi)$  and  $(\tilde{m}, \tilde{\phi})$  two weak solutions of (3.5) with the same initial datum. Set  $\bar{m} = m - \tilde{m}$ ,  $\bar{\phi} = \phi - \tilde{\phi}$  and  $W = |\bar{m}| + |\bar{\phi}|$ . To keep the notation simple, for  $(s, x) \in \mathbb{R}_+ \times \mathbb{R}^d$ , we shall denote  $\bar{m}_s(x) = \bar{m}(s, x)$  and  $\bar{\phi}_s = \bar{\phi}(s, x)$ . For  $1 \leq k \leq d$  and  $s > 0$ , let  $M_k$  and  $F_k$  be defined by

$$\begin{aligned} M_k(m_s, \phi_s) &= (\phi_s - m_s^2) (\partial_k J * m_s), \\ F_k(m_s, \phi_s) &= m_s (1 - \phi_s) (\partial_k J * m_s). \end{aligned}$$

Observe that for all  $s \in [0, t]$ ,

$$\begin{aligned} M_k(m_s, \phi_s) - M_k(\tilde{m}_s, \tilde{\phi}_s) &= (\partial_k J * \bar{m}_s) (\bar{\phi}_s - \tilde{m}_s^2) \\ &\quad + (\partial_k J * m_s) [\bar{\phi}_s - \bar{m}_s (m_s + \tilde{m}_s)] \end{aligned}$$

and

$$\begin{aligned} F_k(m_s, \phi_s) - F_k(\tilde{m}_s, \tilde{\phi}_s) &= (\partial_k J * \bar{m}_s) m_s (1 - \phi_s) \\ &\quad + (\partial_k J * \tilde{m}_s) [\bar{m}_s (1 - \phi_s) - \tilde{m}_s \bar{\phi}_s]. \end{aligned}$$

It follows that there exists a positive constant  $C_2$  that depends on  $\|m\|_\infty$ ,  $\|\phi\|_\infty$  and  $\sup_{1 \leq k \leq d} \|\partial_k J\|$  such that, for almost every  $(s, x) \in [0, t] \times \mathbb{R}^d$ ,

$$\begin{aligned} |M_k(m_s, \phi_s) - M_k(\tilde{m}_s, \tilde{\phi}_s)| &\leq C_2 R(t), \\ |F_k(m_s, \phi_s) - F_k(\tilde{m}_s, \tilde{\phi}_s)| &\leq C_2 R(t). \end{aligned}$$

Here  $R(t)$  stands for the essential sup of  $W$  in  $[0, t] \times \mathbb{R}^d$ :

$$R(t) = \text{ess sup}_{[0,t] \times \mathbb{R}^d} (W(s, x)).$$

Since  $(m, \phi)$  and  $(\tilde{m}, \tilde{\phi})$  are two weak solutions of (3.5), we obtain by (5.1) that for all  $0 \leq \tau \leq t$

$$\begin{aligned} \left| \langle \bar{m}(\tau, \cdot), \mathcal{H}_{\tau,\varepsilon}^f(\tau, \cdot) \rangle \right| &= \beta \left| \sum_{k=1}^d \int_0^\tau \left\langle \left( M_k(m_s, \phi_s) - M_k(\tilde{m}_s, \tilde{\phi}_s) \right), \partial_k \mathcal{H}_{\tau,\varepsilon}^f(s, \cdot) \right\rangle ds \right|, \\ &\leq C_3 \sqrt{t} R(t) \|f\|_1, \\ \left| \langle \bar{\phi}(\tau, \cdot), \mathcal{H}_{\tau,\varepsilon}^f(\tau, \cdot) \rangle \right| &= \beta \left| \sum_{k=1}^d \int_0^\tau \left\langle \left( F_k(m_s, \phi_s) - F_k(\tilde{m}_s, \tilde{\phi}_s) \right), \partial_k \mathcal{H}_{\tau,\varepsilon}^f(s, \cdot) \right\rangle ds \right| \\ &\leq C_3 \sqrt{t} R(t) \|f\|_1, \end{aligned}$$

for some positive constant  $C_3$ . By observing that  $h_\varepsilon$  is an approximate identity in  $\varepsilon$ , we obtain that

$$\begin{aligned} \left| \langle \overline{m}(\tau, \cdot), f \rangle \right| &\leq C_3 \sqrt{t} R(t) \|f\|_1, \\ \left| \langle \overline{\phi}(\tau, \cdot), f \rangle \right| &\leq C_3 \sqrt{t} R(t) \|f\|_1 \end{aligned}$$

for all  $f \in \mathcal{C}_K(\mathbb{R}^d)$  and then for all  $f \in L^1(\mathbb{R}^d)$ . It follows that, for  $0 \leq \tau \leq t$  and  $f \in L^1(\mathbb{R}^d)$ ,

$$\begin{aligned} \langle |\overline{m}(\tau, \cdot)|, f \rangle &\leq C_3 \sqrt{t} R(t) \|f\|_1, \\ \langle |\overline{\phi}(\tau, \cdot)|, f \rangle &\leq C_3 \sqrt{t} R(t) \|f\|_1. \end{aligned}$$

Therefore, for all  $0 \leq \tau \leq t$  and  $f \in L^1(\mathbb{R}^d)$ ,

$$\langle W(\tau, \cdot), f \rangle \leq 2C_3 \sqrt{t} R(t) \|f\|_1, \quad (5.2)$$

which implies that, for all  $\tau \in [0, t]$  (see Appendix),

$$W(\tau, \cdot) \in L^\infty(\mathbb{R}^d) \text{ and } \|W(\tau, \cdot)\|_\infty \leq 2C_3 \sqrt{t} R(t). \quad (5.3)$$

On the other hand, proceeding as in the proof of (5.2), we obtain

$$\langle W(\tau, \cdot), f \rangle \leq 2C_3 \sqrt{t} \tilde{R}(t),$$

where  $\tilde{R}(t)$  is given by

$$\tilde{R}(t) = \sup_{0 \leq s \leq t} \|W(s, \cdot)\|_\infty.$$

This implies that

$$\tilde{R}(t) \leq 2C_3 \sqrt{t} \tilde{R}(t).$$

Choosing  $t = t_0$  such that  $2C_3 \sqrt{t_0} < 1$ , this gives uniqueness in  $[0, t_0] \times \mathbb{R}^d$ . To conclude the proof we have just to repeat the same arguments in  $[t_0, 2t_0]$ , and in each interval  $[kt_0, (k+1)t_0]$ ,  $k \in \mathbb{N}$ ,  $k > 1$ .  $\square$

*Glauber dynamics.*

The proof of the hydrodynamical limit in the Glauber case is based on martingales arguments and does not require Dirichlet form estimates. Following the same strategy as in the Kawasaki case we provide the analogous of Lemma 5.1, 5.2 and 5.4. Fixed a parameter  $\beta > 0$  and the time interval  $[0, T]$ , we denote by  $\mathbf{P}_{\mu^\gamma}^{(G)}$  the law of the process  $(\sigma(t, \cdot))_{t \in [0, T]}$  (without acceleration) on the space  $D([0, T], \Omega)$  and by  $Q_{\mu^\gamma}^{(G)}$  the law of the process  $(\pi_t^{1, \gamma}, \pi_t^{2, \gamma})_{t \in [0, T]}$  on the space  $D([0, T], \mathcal{M}^2)$  with initial distribution  $\mu^\gamma$ . As in the Kawasaki case the proof of the analogous of Lemma 5.1 is simple and is omitted. We shall give only the proof of the analogous of Lemma 5.2: All limit points  $Q^*$  of the sequence  $(Q_{\mu^\gamma}^{(G)})_\gamma$  are concentrated on weak solutions of equation (3.7). Finally the proof of the

uniqueness of the equation (3.7) goes on along the same lines as in the Kawasaki case. It is easier, since we do not need to use properties of the heat kernel.

Denote by  $\mathcal{C}_K^{1,0}([0, T] \times \mathbb{R}^d)$  the space of continuous functions with compact support and with derivative continuous in time. Let  $U = (U^1, U^2) \in \mathcal{C}_K^{1,0}([0, T] \times \mathbb{R}^d) \times \mathcal{C}_K^{1,0}([0, T] \times \mathbb{R}^d)$ , and consider the martingale  $M_t^{(G),U}$  defined by

$$M_t^{(G),U} = \sum_{\iota=1}^2 \left\{ \langle \pi_t^{\iota,\gamma}, U_t^\iota \rangle - \langle \pi_0^{\iota,\gamma}, U_0^\iota \rangle - \int_0^t \left( \partial_s + \mathbb{L}_\gamma^{G,\beta} \right) \langle \pi_s^{\iota,\gamma}, U_s^\iota \rangle ds \right\}.$$

Observe that since for all  $\sigma \in \Omega$  and  $i \in \mathbb{Z}^d$ ,  $\sigma(i) \in \{-1, 0, 1\}$  we have  $1_{\{\sigma(i)=-1\}} = (1/2)\sigma(i)(\sigma(i) - 1)$ ,  $1_{\{\sigma(i)=0\}} = (1 - (\sigma(i))^2)$  and  $1_{\{\sigma(i)=1\}} = (1/2)\sigma(i)(\sigma(i) + 1)$  we may thus rewrite the generator  $\mathbb{L}_\gamma^{(G),\beta}$  as

$$\mathbb{L}_\gamma^{(G),\beta} = \mathbb{L}^{(f)} + \mathbb{L}^- + \mathbb{L}^+,$$

with

$$\begin{aligned} (\mathbb{L}^{(f)}g)(\sigma) &= \frac{1}{2} \sum_{i \in \mathbb{Z}^d} \frac{e^{-\frac{\beta}{2}(\nabla_i^{(f)}H_\gamma)(\sigma)}}{2 \cosh\left(\frac{\beta}{2}(\nabla_i^{(f)}H_\gamma)\right)} \left[ (\nabla_i^{(f)}g)(\sigma) \right], \\ (\mathbb{L}^\mp g)(\sigma) &= \frac{1}{2} \sum_{i \in \mathbb{Z}^d} \frac{e^{-\frac{\beta}{2}(\nabla_i^\mp H_\gamma)(\sigma)}}{2 \cos \cosh\left(\frac{\beta}{2}(\nabla_i^\mp H_\gamma)\right)} \left( (1 - \sigma(i)^2) + \frac{\sigma(i)(\sigma(i) \pm 1)}{2} \right) \left[ (\nabla_i^\mp g)(\sigma) \right], \end{aligned}$$

where for a cylinder function  $F$ ,  $(\nabla_i^{(f)}F)(\sigma)$  is defined by

$$(\nabla_i^{(f)}F)(\sigma) = F(\sigma^{(f),i}) - F(\sigma).$$

For  $i \in \mathbb{Z}^d$ ,  $\sigma^{(f),i}$  is a configuration obtained from  $\sigma$  by flipping the value at  $i$

$$\left( \sigma^{(f),i} \right) (l) = \begin{cases} \sigma(l) & \text{if } l \neq i \\ -\sigma(i) & \text{if } l = i. \end{cases}$$

On the other hand, for all  $\sigma \in \Omega$  and  $i \in \mathbb{Z}^d$  we have

$$\begin{aligned} \exp\left(-\frac{\beta}{2}(\nabla_i^{(f)}H_\gamma)(\sigma)\right) &= \exp\left(-\beta\sigma(i)\alpha^{h_1}(\gamma i)\right) \exp\left(2\gamma^d J_\gamma(0)\right), \\ \exp\left(-\frac{\beta}{2}(\nabla_i^\mp H_\gamma)(\sigma)\right) &= \exp\left(\mp \frac{\beta}{2}[\alpha^{h_1}(\gamma i) + \tilde{h}_2^\gamma(2\sigma(i) \mp 1)]\right) \exp\left(-\gamma^d J_\gamma(0)\right), \end{aligned}$$

where  $\alpha^{h_1}$  and  $\tilde{h}_2^\gamma$  are defined by

$$\alpha^{h_1}(\gamma i) = 2(\pi^{1,\gamma} * J)(\gamma i) + h_1, \quad \tilde{h}_2^\gamma = \left( h_2 - \gamma^d \sum_{k \in \mathbb{Z}^d} J(\gamma k) \right),$$



for  $s \in [0, T]$ , we shall denote

$$\alpha_s^{h_1}(\gamma i) = 2(\pi_s^{1,\gamma} * J)(\gamma i) + h_1.$$

By using the relation  $e^b = (\cosh(b) + \sinh(b))$  it is easy to show that the martingale  $M_t^{(G),U}$  can be written as

$$M_t^{(G),U} = M_t^1 + M_t^2 + o_\gamma(1),$$

where  $M_t^1$  and  $M_t^2$  are the martingales given by

$$\begin{aligned} M_t^1 &= \langle \pi_t^{1,\gamma}, U_t^1 \rangle - \langle \pi_0^{1,\gamma}, U_0^1 \rangle - \int_0^t \langle \pi_s^{1,\gamma}, \partial_s U_s^1 \rangle ds \\ &\quad - \frac{1}{2} \int_0^t \left\{ \langle -\pi_s^{1,\gamma}, U_s^1 \rangle + \langle \pi_s^{2,\gamma}, U_s^1(\cdot) \tanh(\beta \alpha_s^{h_1}(\cdot)) \rangle \right\} ds \\ &\quad - \frac{1}{4} \gamma^d \sum_{i \in \mathbb{Z}^d} \int_0^t U_s^1(\gamma i) \left\{ \tanh\left(\frac{\beta}{2} [\alpha_s^{h_1}(\gamma i) + \tilde{h}_2^\gamma]\right) + \tanh\left(\frac{\beta}{2} [\alpha_s^{h_1}(\gamma i) - \tilde{h}_2^\gamma]\right) \right\} ds \\ &\quad - \frac{1}{8} \int_0^t \left\langle \pi_s^{1,\gamma}, U_s^1(\cdot) \left\{ \tanh\left(\frac{\beta}{2} [\alpha_s^{h_1}(\cdot) + \tilde{h}_2^\gamma]\right) - \tanh\left(\frac{\beta}{2} [\alpha_s^{h_1}(\cdot) - \tilde{h}_2^\gamma]\right) - 2 \right\} \right\rangle ds \\ &\quad - \frac{1}{8} \int_0^t \left\langle -\pi_s^{2,\gamma}, U_s^1(\cdot) \left\{ \tanh\left(\frac{\beta}{2} [\alpha_s^{h_1}(\cdot) + \tilde{h}_2^\gamma]\right) + \tanh\left(\frac{\beta}{2} [\alpha_s^{h_1}(\cdot) - \tilde{h}_2^\gamma]\right) \right\} \right\rangle ds \end{aligned}$$

and

$$\begin{aligned} M_t^2 &= \langle \pi_t^{2,\gamma}, U_t^2 \rangle - \langle \pi_0^{2,\gamma}, U_0^2 \rangle - \int_0^t \langle \pi_s^{2,\gamma}, \partial_s U_s^2 \rangle ds \\ &\quad - \frac{1}{4} \gamma^d \sum_{i \in \mathbb{Z}^d} \int_0^t U_s^2(\gamma i) \left\{ \tanh\left(\frac{\beta}{2} [\alpha_s^{h_1}(\gamma i) + \tilde{h}_2^\gamma]\right) - \tanh\left(\frac{\beta}{2} [\alpha_s^{h_1}(\gamma i) - \tilde{h}_2^\gamma]\right) + 2 \right\} ds \\ &\quad - \frac{1}{8} \int_0^t \left\langle -\pi_s^{2,\gamma}, U_s^2(\cdot) \left\{ \tanh\left(\frac{\beta}{2} [\alpha_s^{h_1}(\cdot) + \tilde{h}_2^\gamma]\right) - \tanh\left(\frac{\beta}{2} [\alpha_s^{h_1}(\cdot) - \tilde{h}_2^\gamma]\right) + 6 \right\} \right\rangle ds \\ &\quad - \frac{1}{8} \int_0^t \left\langle \pi_s^{1,\gamma}, U_s^2(\cdot) \left\{ \tanh\left(\frac{\beta}{2} [\alpha_s^{h_1}(\cdot) + \tilde{h}_2^\gamma]\right) + \tanh\left(\frac{\beta}{2} [\alpha_s^{h_1}(\cdot) - \tilde{h}_2^\gamma]\right) \right\} \right\rangle ds. \end{aligned}$$

On the other hand, a simple computation shows that the quadratic variation  $\mathcal{N}^{(G),1}$  (resp.  $\mathcal{N}^{(G),2}$ ) of the martingale  $M_t^1$  (resp.  $M_t^2$ ) vanishes as  $\gamma \rightarrow 0$ . Therefore, using Chebychev's inequality and Doob's inequality, we obtain

$$\lim_{\gamma \rightarrow 0} \mathbf{P}_{\mu^\gamma}^{(G)} \left\{ \left[ \sup_{0 \leq s \leq T} |M_s^1| + \sup_{0 \leq s \leq T} |M_s^2| \right] > \delta \right\} = 0$$

for any positive  $\delta$ .

To conclude the proof we have just to let  $\gamma \rightarrow 0$  and to follow the same arguments as in the Kawasaki case.  $\square$

## 6. Non gradient dynamics.

In this section, we consider a different kind of dynamics reversible for the Gibbs measure associated to the Hamiltonian (1.1) (with  $h_2 = 1$ ) which is of the so called non gradient type [KL]. We consider a system of  $N$  spins on a  $d$ -dimensional torus  $\mathbb{T}^d$ . At times exponentially distributed each bond  $(i, j) \in \mathbb{T}^d \times \mathbb{T}^d, |i - j| = 1$  change its configuration  $(\sigma(i), \sigma(j))$  independently of the others (or stays unchanged) to the new configuration  $(\sigma'(i), \sigma'(j))$  in such a way that  $|\sigma(i) - \sigma'(i)| = 1, |\sigma(j) - \sigma'(j)| = 1$  and  $\sigma(i) + \sigma(j) = \sigma'(i) + \sigma'(j)$  with jump rates chosen to satisfy the detailed balance condition with respect the Hamiltonian

$$H_\gamma(\eta) = - \sum_{i,j \in \mathbb{T}^d} J_\gamma(i-j) \sigma(i) \sigma(j),$$

In other words, the transitions allowed for a bond  $(i, j)$  are

$$\begin{aligned} (0, -1) &\iff (-1, 0), & (1, 0) &\iff (0, 1) \\ (1, -1) &\iff (0, 0), & (0, 0) &\iff (-1, 1) \end{aligned}$$

We remark that the difference between the number of positive and negative spins is conserved by this dynamics, while the number of zero spins is not, because negative and positive neighbouring spins can annihilate to create two spins with zero value or viceversa two zero spins can disappear to give rise to a couple of spins  $\pm 1$ .

This dynamics, when reformulated as a lattice gas, turns out to be at  $\beta = 0$  the generalized exclusion process introduced in [KLO]. To match the notations in that paper we prefer to use in this section the representation of the system in terms of the occupation number  $\eta(i) = 0, 1, 2$  instead of the spin variable  $\sigma(i) = -1, 0, 1$ , their relation being  $\sigma(i) = \eta(i) - 1$ . In each site of the torus  $\mathbb{T}^d$  there are at most 2 particles. A configuration of the system is an element  $\eta$  of  $\mathbb{X}_N = \{0, 1, 2\}^{\mathbb{T}^d}$ , where  $N$  is the number of sites in  $\mathbb{T}^d$ . Particles move on the torus in the following way. A particle in  $i$  jumps with a given rate to the nearest neighbour  $j$  if in  $j$  there is at most one particle. We call  $\eta^{i,j}$  the configuration obtained from  $\eta$  letting one particle jump from  $i$  to  $j$ :

$$(\eta^{i,j})(k) = \begin{cases} \eta(k) & \text{if } k \neq i, j, \\ \eta(k) - 1 & \text{if } k = i, \\ \eta(k) + 1 & \text{if } k = j. \end{cases}$$

For  $(i, j) \in \mathbb{T}^d$  and every cylindric function  $F : \mathbb{X}_N \rightarrow \mathbb{R}$ , define  $(\nabla_{i,j} F)(\eta)$  by

$$(\nabla_{i,j} F)(\eta) = r_{i,j}(\eta) \{F(\eta^{i,j}) - F(\eta)\}.$$

where

$$r_{i,j}(\eta) = 1\{\eta(i) > 0, \eta(j) < 2\},$$

The jump rates are

$$C_\gamma^\beta(i, j; \eta) = \Phi \{ \beta (H_\gamma(\eta^{i,j}) - H_\gamma(\eta)) \},$$

with

$$H_\gamma(\eta) = - \sum_{i,j \in \mathbb{T}^d} J_\gamma(i-j) \eta(i) \eta(j), \quad (6.1)$$

where  $J_\gamma$  is the Kac potential defined in section 2 and  $\Phi : \mathbb{R} \rightarrow \mathbb{R}_+$  is a continuously differentiable function in a neighborhood of 0, such that  $\Phi(0) = 1$ , satisfying the detailed balance condition

$$\Phi(E) = \exp(-E) \Phi(-E). \quad (6.2)$$

The generator of this jump Markov process  $(\eta_t)_{t \geq 0}$  is given by

$$\left( \mathbb{L}_\gamma^\beta f \right) (\eta) = (1/2) \sum_{\substack{i,j \in \mathbb{T}^d \\ |i-j|=1}} C_\gamma^\beta(i, j; \eta) \{ (\nabla_{i,j} f) (\eta) \}.$$

where we have made explicit the dependence on the parameter  $\beta \geq 0$ .

Lemma 6.9 shows that the dynamics with parameter  $\beta > 0$  is a weak perturbation of the generalized simple exclusion process GSEP in [KLO] and reduces to it at  $\beta = 0$ . We shall denote the generator of GSEP by  $\mathbb{L}_\gamma^0$ .

For  $\varphi \geq 0$ , define  $\bar{\nu}_\varphi^N$  as the product measure on  $\mathbb{X}_N$  with marginals given by

$$\bar{\nu}_\varphi^N \{ \eta(0) = r \} = \frac{\varphi^r}{1 + \varphi + \varphi^2}, \quad r = 0, 1, 2.$$

Let  $R(\varphi)$  be the mean occupation number of particles under  $\bar{\nu}_\varphi^N$  :

$$R(\varphi) = E_{\bar{\nu}_\varphi^N} [\eta(0)].$$

The function  $R : \mathbb{R}_+ \rightarrow [0, 2)$  is a bijection and we denote by  $\psi : [0, 2) \rightarrow \mathbb{R}_+$  its inverse. For every  $\alpha$  in  $[0, 2)$ , we denote by  $\nu_\alpha^N$  the product measure  $\bar{\nu}_{\psi(\alpha)}^N$  so that the density of particles on each site is  $\alpha$  :

$$E_{\nu_\alpha^N} [\eta(x)] = \alpha \quad \text{for } x \text{ in } \mathbb{X}_N.$$

We will use the notation  $\nu_\alpha$  for the product measure on the infinite volume product space  $\mathbb{X} = \{0, 1, 2\}^{\mathbb{Z}^d}$  and  $\langle f \rangle_\alpha$  for the expectation of a cylinder function  $f$  with respect to  $\nu_\alpha$  or  $\nu_\alpha^N$  :

$$\langle f \rangle_\alpha = \int f(\eta) \nu_\alpha(d\eta). \quad (6.3)$$

The one-parameter family  $(\nu_\alpha^N)_\alpha$  of probability measures is reversible for the generator  $\mathbb{L}_\gamma^0$  (GSEP) and for  $\beta > 0$  the one-parameter family of probability measures  $(\nu_\alpha^{\beta,N})_\alpha$  given by

$$\nu_\alpha^{\beta,N}(\eta) = \frac{\exp\{-\beta H_\gamma(\eta)\}}{Z_\alpha^\beta} \nu_\alpha^N(\eta)$$

is reversible for the dynamics with  $\beta > 0$ . Here  $Z_\alpha^\beta$  is the normalization constant.

We now choose  $N = \gamma^{-1}$  and speed up the generator as  $\gamma^{-2}$ , as in the Kawasaki case, and study the limit  $N \rightarrow \infty$ . In this Section we show that, starting from a sequence of measures on  $\mathbb{X}_N$  associated to the same initial profile  $\rho_0$ , the density field converges, as  $N$  increases to infinity, to the weak solution of the nonlinear parabolic equation (6.5), where the diffusion matrix  $D$  is given [KLO] by (6.4) below.

It has been proved in [KLO] that the coefficients  $D_{k,m}(\rho)$  are nonlinear continuous functions of  $\rho$  and that  $D$  is strictly elliptic. That is not enough to prove the uniqueness of weak solutions of (6.5), which is easy to prove instead if the diffusion coefficient is known to be locally Lipschitz continuous (for example by the method in [LMS]).

In order to define the diffusion coefficient, we need to establish some notation and to consider the generalized exclusion process in the infinite volume space  $\mathbb{X}$ .

For  $i$  in  $\mathbb{Z}^d$ , let  $\tau_i$  denote the space shift by  $i$  units on  $\mathbb{X}$ . For a cylinder function  $F$  on  $\mathbb{X}$ , define the formal sum

$$\Gamma_F(\eta) = \sum_{j \in \mathbb{Z}^d} (\tau_j F)(\eta)$$

which does not make sense but for which the quantities  $\{\nabla_{0,e_k} \Gamma_F, 1 \leq k \leq d\}$  are well defined. Here  $\{e_1, \dots, e_d\}$  are the unitary vectors in the coordinate directions of  $\mathbb{Z}^d$ . For each  $\alpha$  in  $[0, 2]$ , let  $D(\alpha) = \{D_{k,m}(\alpha), 1 \leq k, m \leq d\}$  be the symmetric matrix defined by the following variational formula

$$a \cdot D(\alpha) a = \frac{1}{2\chi(\alpha)} \inf_F \sum_{k=1}^d \left\langle \left( a_k r_{0,e_k} + \nabla_{0,e_k} \Gamma_F \right)^2 \right\rangle_\alpha, \quad (6.4)$$

for any vector  $a$  in  $\mathbb{R}^d$ . In this formula  $\chi(\alpha)$  is the static compressibility defined by

$$\chi(\alpha) = \langle \eta(0)^2 \rangle_\alpha - \langle \eta(0) \rangle_\alpha^2 = \langle \sigma(0)^2 \rangle_\alpha - \langle \sigma(0) \rangle_\alpha^2.$$

For a measure  $\mu$  on  $\mathbb{X}_N$ , denote by  $P_\mu$  the probability measure on the path space  $D(\mathbb{R}_+, \mathbb{X}_N)$  corresponding to the Markov process  $(\eta_t)$  with generator speeded up by  $N^2$  and starting from  $\mu$ , and by  $\mathbf{E}_{P_\mu}$  the expectation with respect to  $P_\mu$ .

Let  $\mathcal{M} = \mathcal{M}(\mathbb{T}^d)$  be the space of positive measures on the  $d$ -dimensional torus  $\mathbb{T}^d$  with total mass bounded by  $2d$ . For each configuration  $\eta$ , denote by  $\pi^N = \pi^N(\eta)$  the positive measure obtained assigning mass  $N^{-d}$  to each particle of  $\eta$ :

$$\pi^N = N^{-d} \sum_{j \in \mathbb{T}^d} \eta(j) \delta_{j/N},$$

where  $\delta_x$  is the Dirac measure concentrated on  $x$ . For each  $t \geq 0$ , denote by  $\pi_t = \pi_t^N$  the empirical measure at time  $t$  :  $\pi_t = \pi^N(\eta_t)$ . For a continuous function  $U$  and  $\pi$  in  $\mathcal{M}(\mathbf{T}^d)$ , we shall denote by  $\langle \pi, U \rangle$  the integral of the function  $U$  with respect to the measure  $\pi$ .

Fix  $T > 0$ . For each probability measure  $\mu$  on  $\mathbb{X}_N$ , denote by  $Q_\mu^N$  the measure on the state space  $D([0, T], \mathcal{M})$  induced by the Markov process  $\pi_t$  speeded up by  $N^2$  and  $\mu^N$ .

**Theorem 6.1** *Consider a sequence of probability measures  $\mu^N$  on  $\mathbb{X}_N$  associated to the initial profile  $\rho_0$  in the following sense :*

$$\lim_{N \rightarrow \infty} \mu^N \left\{ \left| \langle \pi^N(\eta), U \rangle - \langle \rho_0(x) dx, U \rangle \right| > \delta \right\} = 0$$

for every continuous function  $U : \mathbf{T}^d \rightarrow \mathbb{R}$  and every  $\delta > 0$ . Then, the sequence of probability measures  $\{Q_\mu^N, N > 1\}$  is tight and all its limit points  $Q^*$  are concentrated on absolutely continuous paths  $\pi(t, dx) = \rho(t, x)dx$  whose density  $\rho$  is the weak solution of the equation

$$\begin{cases} \partial_t \rho = \sum_{k,m=1}^d \partial_k \left\{ D_{k,m}(\rho) \left\{ \partial_m \rho - 2\beta \chi(\rho) (\partial_m J * \rho) \right\} \right\}, \\ \rho(0, \cdot) = \rho_0(\cdot). \end{cases} \quad (6.5)$$

and belongs to  $L^2([0, T], H_1(\mathbf{T}^d))$ . Moreover if the diffusion matrix  $D$  is locally Lipschitz continuous, then the empirical measures converge in the limit  $N \rightarrow \infty$  to the unique weak solution of equation (6.5).

Since  $\rho(x) = m(x) + 1$  the equation for the magnetization is the same as (6.5). It is not difficult to see that it can be put in the form

$$\partial_t m = \nabla \cdot \left( \Sigma \nabla \frac{\delta \mathcal{G}}{\delta m} \right) \quad (6.6)$$

with the energy functional

$$\mathcal{G}(m(r)) := \int dr g^0(m(r)) + \frac{1}{2} \int dr \int dr' J(r - r') m(r) m(r') \quad (6.7)$$

where

$$g^0(m) := -m^2 - \beta^{-1} s(m, \phi(m)) \quad (6.8)$$

and  $\phi(m) = \langle \sigma^2 \rangle_m$ , where  $\langle \cdot \rangle_m$  is defined in (6.3). The mobility is given by the Einstein relation  $\Sigma = D(m)\chi(m)$ . Moreover the energy functional  $\mathcal{G}$  is a Liapunov functional for (6.5).

The stationary homogeneous solutions of (6.6) are given by the solutions of (2.8) with  $h_1 = 0$  and  $h_2 = 1$ . Notice that for these values of the parameters the second equation of (2.8) determines the function  $\phi(m)$  defined above as

$$\phi(m) = \frac{\cosh 2\beta m}{1 + \cosh 2\beta m}.$$

For  $\beta > 3/4$  the function  $g^0(m)$  has two symmetric minima  $\pm m_s$  determined by the non vanishing solutions of

$$m = \frac{\sinh 2\beta m}{1 + \sinh 2\beta m}$$

**Proof of Theorem 6.1.**

Following the strategy adopted in Section 4 for the Kawasaki dynamics, we divide the proof of Theorem 6.1 in three steps: Tightness, identification of the limit, and under the assumption that the diffusion matrix is locally Lipschitz continuous, the uniqueness of the hydrodynamic equation.

The proof of the tightness is essentially the same as the one given in Section 6 in [LMS] and therefore is omitted. Notice however that in the present case of perturbed generalized simple exclusion process the invariant measures are not product while the proof of tightness in [KLO] use explicitly the fact that the product measures  $(\nu_\alpha^N)_\alpha$  are invariant.

For the uniqueness of weak solutions of the hydrodynamic equation we need an energy estimate which states that every limit point  $Q^*$  of the sequence  $\{Q_\mu^N, N > 1\}$  is concentrated on paths whose density  $\rho$  belongs to  $L^2([0, T], H_1(\mathbf{T}^d))$ .

**Proposition 6.2** *Let  $Q^*$  be a limit point of the sequence  $\{Q_\mu^N, N > 1\}$ . Then,*

$$E_{Q^*} \left[ \int_0^T ds \left( \int_{\mathbf{T}^d} \|\nabla \rho(s, x)\|^2 dx \right) \right] < \infty .$$

The proof of this proposition is similar to the proof of Proposition A.1.1. in [KLO] and is therefore omitted. The proof of the uniqueness of weak solutions follows the same lines as in [VY] or in [LMS]. It is here that we need the diffusion coefficient to be locally Lipschitz.

The identification of the limit is not trivial and is the main step of the proof of Theorem 6.1.

**Lemma 6.3** *All limit points  $Q^*$  of the sequence  $\{Q_\mu^N, N > 1\}$  are concentrated on the path  $\rho(t, x)dx$  whose density  $\rho$  is the weak solution of equation (6.5).*

*Proof.* Fix a function  $U$  in  $\mathcal{C}^{1,2}([0, T] \times \mathbf{T}^d)$ . Consider the martingales  $M_t^U = M_t^{U, N}$ ,

$N_t^U = N_t^{U,N}$  defined by

$$\begin{aligned} M_t^U &= \langle \pi_t^N, U_t \rangle - \langle \pi_0^N, U_0 \rangle - \int_0^t (\partial_s + N^2 \mathbb{L}_\gamma^\beta) \langle \pi_s^N, U_s \rangle ds, \\ N_t^U &= (M_t^U)^2 \\ &\quad - \int_0^t \left\{ N^2 \mathbb{L}_\gamma^\beta (\langle \pi_s^N, U_s \rangle)^2 - 2 \langle \pi_s^N, U_s \rangle N^2 \mathbb{L}_\gamma^\beta \langle \pi_s^N, U_s \rangle \right\} ds. \end{aligned}$$

In these formulas, for a continuous function  $U$  and  $\pi$  in  $\mathcal{M}(\mathbb{T}^d)$ ,  $\langle \pi, U \rangle$  stands for the integral of the function  $U$  with respect to the measure  $\pi$ .

A simple computation of the integral term of  $N_t^U$  shows that the expectation of the quadratic variation of the martingale  $M_t^U$  vanishes as  $N \uparrow \infty$ . Therefore, by Doob's inequality, for every  $\delta > 0$ ,

$$\lim_{N \rightarrow \infty} P_{\mu^N} \left[ \sup_{0 \leq t \leq T} |M_t^U| > \delta \right] = 0. \quad (6.9)$$

We now turn to the martingale  $M_t^U$ . A summation by parts permits to rewrite the integral term of the martingale  $M_t^U$  as

$$\int_0^t \langle \pi_s^N, \partial_s U_s \rangle ds + \int_0^t ds N^{-d+1} \sum_{k=1}^d \sum_{i \in \mathbb{T}^d} (\partial_k^N U)(s, i/N) \mathbf{W}_{i, i+e_k}^{\beta, N}(\eta_s),$$

where  $\mathbf{W}_{i, i+e_k}^{\beta, N}(\eta)$  is the current over the bond  $\{i, i+e_k\}$  :

$$\mathbf{W}_{i, i+e_k}^{\beta, N}(\eta) = \frac{1}{2} \{ C_\gamma^\beta(i, i+e_k; \eta) - C_\gamma^\beta(i+e_k, i; \eta) \}$$

and  $(\partial_k^N U)$  is the discrete gradient defined by

$$(\partial_k^N U)(i/N) = N[U((i+e_k)/N) - U(i/N)].$$

Remark that we may write the current  $\mathbf{W}_{i, i+e_k}^{\beta, N}$  as the sum of the current  $W_{i, i+e_k}$  of the GSEP ( $\mathbb{L}_\gamma^0$ ) and a term coming from the perturbation:

$$\mathbf{W}_{i, i+e_k}^{\beta, N}(\eta) = W_{i, i+e_k}(\eta) + \frac{\beta}{2} N^{-1} (\partial_k J * \pi^N(\eta))(i/N) \{r_{i, i+e_k} + r_{i+e_k, i}\} + N^{-1} o_N(1),$$

where  $o_N(1)$  is a random variable which is bounded in absolute value by a constant that converges to 0 as  $N \uparrow \infty$  and

$$W_{i, i+e_k}(\eta) = \frac{1}{2} \{r_{i, i+e_k}(\eta) - r_{i+e_k, i}(\eta)\}.$$

We will often omit the dependence of  $W_{i,i+e_k}$  and  $\mathbf{W}_{i,i+e_k}^{\beta,N}$  on  $\eta$  and  $N$ .

Following the non gradient method of Varadhan [V] and the entropy method of [GPV] we now replace the current  $\mathbf{W}_{i,i+e_k}^{\beta,N}$  appearing in the integral term of the martingale  $M_t^U$  by a linear combination of the gradient  $\left\{ \eta(i+e_m) - \eta(i) \right\}$  and  $\left\{ \beta N^{-1} (\partial_m J * \pi^N(\eta)) (i/N) \right\}$ . This requires some notations. For  $\ell \leq N$  and  $i$  in  $\mathbf{T}^d$ , let  $\eta^\ell(x)$  stands for the mean number of particles in a cube of size  $2\ell + 1$  centered in  $i$  :

$$\eta^\ell(i) = \frac{1}{(2\ell + 1)^d} \sum_{|j-i| \leq \ell} \eta(j) .$$

For  $k = 1, \dots, d$ ,  $N \geq 1$ ,  $\varepsilon > 0$  and a smooth function  $G : \mathbf{T}^d \rightarrow \mathbb{R}$ , let

$$X_{N,\varepsilon}^k(G, \eta) = N^{-d+1} \sum_{i \in \mathbf{T}^d} G(i/N) \tau_i \mathbf{W}_k^{N\varepsilon}(\eta) ,$$

where

$$\mathbf{W}_k^{N\varepsilon}(\eta) = \mathbf{W}_{0,e_k} + \sum_{m=1}^d D_{k,m}(\eta^{N\varepsilon}(0)) \left\{ \left[ \eta^{N\varepsilon}(e_m) - \eta^{N\varepsilon}(0) \right] \right. \\ \left. - 2\beta N^{-1} \chi(\eta^{N\varepsilon}(0)) (\partial_m J * \pi^N(\eta)(0)) \right\} .$$

Next theorem is the main step in the proof of Lemma 6.3 and therefore of the hydrodynamic limit.

**Theorem 6.4** *For every smooth function  $G : [0, T] \times \mathbf{T}^d \rightarrow \mathbb{R}$ ,*

$$\limsup_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} E_{P_{\mu^N}} \left[ \left| \int_0^T X_{N,\varepsilon}^k(G_s, \eta_s) ds \right| \right] = 0$$

for  $k = 1, \dots, d$ .

The proof of this theorem is postponed to the end of this section. We show now how Theorem 6.4 allows to conclude the proof of Theorem 6.1. For  $1 \leq k, m \leq d$ , denote by  $d_{k,m}$  the integral of  $D_{k,m}$  :

$$d_{k,m}(\alpha) = \int_0^\alpha D_{k,m}(u) du , \quad \text{for } \alpha \in [0, 2] .$$

Since  $D_{k,m}$  is a continuous function, by Taylor expansion,

$$d_{k,m}(\eta^{N\varepsilon}(e_m)) - d_{k,m}(\eta^{N\varepsilon}(0)) \\ = D_{k,m}(\eta^{N\varepsilon}(0)) \left\{ \eta^{N\varepsilon}(e_m) - \eta^{N\varepsilon}(0) \right\} + (N\varepsilon)^{-1} o_N(1) .$$



It follows therefore from Theorem 6.4 that

$$\limsup_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} E_{P_{\mu^N}} \left[ \left| \int_0^T \mathbb{Y}_{N,\varepsilon}^k(G_s, \eta_s) ds \right| \right] = 0 ,$$

where

$$\mathbb{Y}_{N,\varepsilon}^k(G, \eta) = N^{1-d} \sum_{i \in \mathbb{T}^d} G(i/N) \tau_i \mathbb{W}_k^{N\varepsilon}(\eta)$$

and

$$\begin{aligned} \mathbb{W}_k^{N\varepsilon}(\eta) &= \mathbf{W}_{0,e_k} + \sum_{m=1}^d \left\{ d_{k,m}(\eta^{N\varepsilon}(e_m)) - d_{k,m}(\eta^{N\varepsilon}(0)) \right\} \\ &\quad - 2\beta N^{-1} \sum_{m=1}^d \left\{ \chi(\eta^{N\varepsilon}(0)) D_{k,m}(\eta^{N\varepsilon}(0)) (\partial_m J * \pi^N(\eta))(0) \right\}. \end{aligned}$$

A summation by parts permits to rewrite the second term of  $\mathbb{Y}_{N,\varepsilon}^i(G, \eta)$  as

$$\begin{aligned} &- N^{-d} \sum_{m=1}^d \sum_{i \in \mathbb{T}^d} (\partial_m G)(i/N) d_{k,m}(\eta^{N\varepsilon}(i)) \\ &- N^{-d} \sum_{m=1}^d \sum_{i \in \mathbb{T}^d} G(i/N) (2\beta \chi D_{k,m})(\eta^{N\varepsilon}(i)) (\partial_m J * \pi^N(\eta))(i/N) + O(N^{-1}). \end{aligned}$$

This concludes the proof of the Theorem.

**Proof of Theorem 6.4.** We first introduce some notations and recall some tools used in the non gradient methods. We denote by  $\mathbb{L}^0$  the pregenerator of GSEP in infinite volume and consider the family  $(\nu_\alpha)$  of invariant measures for  $\mathbb{L}^0$ . Let  $\mathcal{C}$  be the space of cylinder functions. For each box  $\Lambda \subset \mathbb{Z}^d$  and a positive integer  $\ell$ , such that  $0 \leq \ell \leq 2|\Lambda|$ , we denote by  $\nu_{\Lambda,\ell}$  the canonical measure on  $\{0, 1, 2\}^\Lambda$  with density  $\ell/|\Lambda|$ . For a cylinder function  $g \in \mathcal{C}$ , denote by  $\Lambda_g$  the smallest rectangle that contains the support of  $g$  and by  $s_g$  the smallest positive integer  $s$  such that  $\Lambda_g \subset \Lambda_s$ . Let  $\mathcal{C}_0$  be the linear space of cylinder functions with mean zero with respect to all canonical invariant measures for  $\mathbb{L}^0$  :

$$\mathcal{C}_0 = \left\{ g \in \mathcal{C} ; \langle g \rangle_{\Lambda_g, \ell} = 0 \text{ for all } 0 \leq \ell \leq 2|\Lambda_g| \right\}.$$

Here  $\langle g \rangle_{\Lambda_g, \ell}$  stands for the expectation of the function  $g$  with respect to the measure  $\nu_{\Lambda_g, \ell}$ .

For a positive density  $0 \leq \rho \leq 2$ , define the semi-norm  $\sqrt{\langle\langle g \rangle\rangle_\rho}$  by the central limit theorem variances

$$\langle\langle g \rangle\rangle_\rho = \lim_{\ell \rightarrow \infty} (2\ell)^{-d} \left\langle \left( -\mathbb{L}_{\Lambda_\ell}^0 \right)^{-1} \sum_{|i| \leq \ell_g} \tau_i g, \sum_{|i| \leq \ell_g} \tau_i g \right\rangle_{\ell, K_\ell},$$

for  $g \in \mathcal{C}_0$  and a sequence of positive integers  $K_\ell$  such that  $0 \leq K_\ell \leq 2(2\ell + 1)^d$  and  $\lim_{\ell \rightarrow \infty} K_\ell / (2\ell)^d = \rho$ . In this formula  $\ell_g = \ell - s_g - 1$ ,  $\Lambda_\ell = \{-\ell, \dots, \ell\}^d$  and for a box  $\Lambda \subset \mathbb{Z}^d$ ,  $\mathbb{L}_\Lambda^0$  is the restriction of  $\mathbb{L}^0$  to the box  $\Lambda$ .

By polarization we may define from  $\ll \cdot \gg_\rho$  a semi-inner product on  $\mathcal{C}_0$ . Moreover, if for cylinder functions  $g$  and  $h$  in  $\mathcal{C}_0$  we define

$$\ll g, h \gg_{\rho,0} = \sum_i \langle g, \tau_i h \rangle_\rho$$

we obtain by the definition of  $\ll \cdot \gg_\rho$  the following properties (cf. [KL]) : For all  $h, g \in \mathcal{C}_0$  and for each  $0 \leq k, m \leq d$

$$\begin{aligned} \ll g, \mathbb{L}^0 h \gg_\rho &= -\ll g, h \gg_{\rho,0}, \\ \ll \eta(e_k) - \eta(0), \mathbb{L}^0 g \gg_\rho &= 0, \\ \ll \eta(e_k) - \eta(0), W_{0,e_m} \gg_\rho &= -\chi(\rho) \delta_{k,m}, \\ \ll W_{0,e_k}, W_{0,e_m} \gg_\rho &= \frac{1}{2} \langle r_{0,e_1} \rangle_\rho \delta_{k,m}, \end{aligned} \tag{6.10}$$

where  $\delta_{k,m}$  is the Kroenecker delta. Let us denote by  $\mathcal{F}$  the space of functions  $F : [0, 2] \times \mathbb{X} \rightarrow \mathbb{R}$  such that

- (i) for each  $\rho \in [0, 2]$ ,  $F(\rho, \cdot)$  is a cylinder function with uniform support i.e. there exists a finite set  $\Lambda \subset \mathbb{Z}^d$  that contains the support of  $F(\rho, \cdot)$  for all  $\rho \in [0, 2]$ .
- (ii) For each configuration  $\eta$ ,  $F(\cdot, \eta)$  is a smooth function.

It has been proved in [KLO] and [KL] (Corollary 7.5.9 in [KL]) that for all  $0 \leq k \leq d$ ,

$$\inf_{F \in \mathcal{F}} \sup_{0 \leq \rho \leq 2} \ll W_{0,e_k} + \sum_{m=1}^d D_{k,m}(\rho) [\eta(e_m) - \eta(0)] - \mathbb{L}^0 F(\rho, \eta) \gg_\rho = 0.$$

For each positive integer  $n \geq 1$  let  $F^{k,n} \in \mathcal{F}$  such that for all  $0 \leq \rho \leq 2$ ,

$$\ll W_{0,e_k} + \sum_{m=1}^d D_{k,m}(\rho) [\eta(e_k) - \eta(0)] - \mathbb{L}^0 F^{k,n}(\rho, \eta) \gg_\rho \leq \frac{1}{n}.$$

It is easy to see that for each  $n \geq 1$  (cf. proof of (7.1.2) in [KL])

$$\limsup_{N \rightarrow \infty} \mathbf{E}_{P_{\mu^N}} \left[ \left| \int_0^T ds N^{1-d} \sum_{i \in \mathbb{T}^d} G(s, i/N) \tau_i \mathbb{L}_\gamma^\beta F^{k,n}(\eta_s^{\varepsilon N}(0), \eta_s) \right| \right] = 0.$$

In particular, to prove the theorem we have to show that

$$\lim_{n \rightarrow \infty} \limsup_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} \mathbf{E}_{P_{\mu^N}} \left[ \left| \int_0^T Z_{N,\varepsilon}^{F^{k,n}}(G_s, \eta_s) ds \right| \right] = 0, \tag{6.11}$$

where  $Z_{N,\varepsilon}^{F^{k,n}}$  ( $G, \eta$ ) is defined by

$$Z_{N,\varepsilon}^{F^{k,n}}(G, \eta) = X_{N,\varepsilon}^k(G, \eta) - N^{1-d} \sum_{i \in \mathbf{T}^d} G(i/N) \tau_i \mathbb{L}_\gamma^\beta F^{k,n}(\eta^{\varepsilon N}(0), \eta).$$

On the other hand, since  $F^{k,n}$  is a cylinder function, it is easy to see by Lemma 6.9 that for all  $i \in \mathbf{T}^d$

$$\begin{aligned} \tau_i \mathbb{L}_\gamma^\beta F^{n,k} &= N^{-1} o_N(1) + \tau_i \mathbb{L}_\gamma^0 F^{n,k} \\ &+ N^{-1} \frac{\beta}{2} \sum_{m=1}^d (\partial_m J * \pi^N(i/N)) \tau_i \left\{ \sum_{j \in \mathbf{T}^d} \left\{ \nabla_{j,j+e_m} F^{n,k} - \nabla_{j+e_m,j} F^{n,k} \right\} \right\}. \end{aligned}$$

We now decompose  $Z_{N,\varepsilon}^{F^{k,n}}$  in two parts,  $Z_{N,\varepsilon}^{F^{k,n},1}$  and  $Z_{N,\varepsilon}^{F^{k,n},2}$  :

$$Z_{N,\varepsilon}^{F^{k,n}} = Z_{N,\varepsilon}^{F^{k,n},1} + Z_{N,\varepsilon}^{F^{k,n},2} + N^{-1} o_N(1),$$

where

$$\begin{aligned} Z_{N,\varepsilon}^{F^{k,n},1} &= N^{1-d} \sum_{i \in \mathbf{T}^d} G(i/N) \tau_i \left\{ W_{0,e_k} + \sum_{m=1}^d D_{k,m}(\eta^{N\varepsilon}(0)) \left[ \eta^{N\varepsilon}(e_m) - \eta^{N\varepsilon}(0) \right] \right\} \\ &- N^{1-d} \sum_{i \in \mathbf{T}^d} G(i/N) \tau_i \mathbb{L}_\gamma^0 F^{n,k}(\eta^{\varepsilon N}(0), \eta), \\ Z_{N,\varepsilon}^{F^{k,n},2} &= \beta N^{-d} \sum_{i \in \mathbf{T}^d} G(i/N) \tau_i \left\{ (\partial_k J * \pi^N(\eta)(0)) \left( \frac{r_{0,e_k} + r_{e_k,0}}{2} \right) \right. \\ &\quad \left. - 2 \sum_{m=1}^d (\chi D_{k,m})(\eta^{N\varepsilon}(0)) (\partial_m J * \pi^N(\eta)(0)) \right\} \\ &- \beta N^{-d} \sum_{i \in \mathbf{T}^d} G(i/N) \tau_i \left\{ \sum_{m=1}^d (\partial_m J * \pi^N(\eta)(0)) \times \right. \\ &\quad \left. \times \left\{ \sum_{j \in \mathbf{T}^d} (1/2) (\nabla_{j,j+e_m} F^{n,k} - \nabla_{j+e_m,j} F^{n,k}) \right\} \right\}. \end{aligned}$$

Here  $(\chi D_{k,m})$  represents the product function  $(\chi D_{k,m})(\alpha) = \chi(\alpha) D_{k,m}(\alpha)$ . To conclude the proof of the theorem it is enough to prove the following lemmas :

**Lemma 6.5** *For each  $0 \leq k \leq d$ ,*

$$\lim_{n \rightarrow \infty} \limsup_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} \mathbf{E}_{P_{\mu^N}} \left[ \left| \int_0^T Z_{N,\varepsilon}^{F^{k,n},1}(G_s, \eta_s) ds \right| \right] = 0. \quad (6.12)$$

**Lemma 6.6** For each  $0 \leq k \leq d$ ,

$$\lim_{n \rightarrow \infty} \limsup_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} \mathbf{E}_{P_{\mu^N}} \left[ \left| \int_0^T Z_{N,\varepsilon}^{F^{k,n},2}(G_s, \eta_s) ds \right| \right] = 0. \quad (6.13)$$

*Proof of Lemma 6.5.* Fix  $0 \leq k \leq d$  and denote  $Z_{N,\varepsilon}^{F^{n,k},1}$  by  $Z_{N,\varepsilon}^1$ . Since the entropy  $s(P_{\mu^N} | P_{\nu_\alpha^N})$  of  $P_{\mu^N}$  with respect to  $P_{\nu_\alpha^N}$  is equal to  $s(\mu^N | \nu_\alpha^N)$  which is bounded by  $CN^d$  for some positive constant  $C$ , by the entropy inequality for any positive  $A$

$$\begin{aligned} \mathbf{E}_{P_{\mu^N}} \left[ \left| \int_0^t Z_{N,\varepsilon}^1(G_s, \eta_s) ds \right| \right] &\leq \frac{C}{A} \\ &+ \frac{1}{AN^d} \log \mathbf{E}_{P_{\nu_\alpha^N}} \left[ \exp \left( AN^d \left| \int_0^t Z_{N,\varepsilon}^1(G_s, \eta_s) ds \right| \right) \right]. \end{aligned}$$

Since  $e^{|x|} \leq e^x + e^{-x}$  and

$$\limsup N^{-d} \log \{a_N + b_N\} \leq \max \{ \limsup N^{-d} \log a_N, \limsup N^{-d} \log b_N \},$$

the absolute value appearing in the exponent of the right hand side of the last inequality can be eliminated. Indeed, by the Feynman-Kac formula, we have

$$\begin{aligned} &\limsup_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} \frac{1}{AN^d} \log \mathbf{E}_{P_{\nu_\alpha^N}} \left[ \exp \left( AN^d \left| \int_0^t Z_{N,\varepsilon}^1(G_s, \eta_s) ds \right| \right) \right] \\ &\leq \limsup_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} \frac{1}{AN^d} \int_0^t \left( \lambda_{N,\varepsilon}(G_s) + \lambda_{N,\varepsilon}(-G_s) \right) ds, \end{aligned}$$

where  $\lambda_{N,\varepsilon}(\pm G_s)$  is the largest eigenvalue of the reversible operator

$$\frac{N^2}{2} \left( \mathbf{L}_\gamma^\beta + \mathbf{L}_\gamma^{\beta,*} \right) + AN^d Z_{N,\varepsilon}^1(\pm G_s, \eta)$$

given by the variational formula

$$\sup_{\substack{f \geq 0 \\ \int f d\nu_\alpha^N = 1}} \left\{ AN^d \int Z_{N,\varepsilon}^1(\pm G_s, \eta) f(\eta) d\nu_\alpha^N(\eta) + N^2 \left\langle \sqrt{f}, \mathbf{L}_\gamma^\beta \sqrt{f} \right\rangle_\alpha \right\}.$$

Here,  $\mathbf{L}_\gamma^{\beta,*}$  stands for the adjoint operator of  $\mathbf{L}_\gamma^\beta$  in  $L^2(\nu_\alpha^N)$ . From Lemma 6.8 below we have, for all positive  $A$

$$\begin{aligned} &\frac{1}{AN^d} \int_0^t \lambda_{N,\varepsilon}(\pm G_s) ds \leq \frac{C_1}{A} t \\ &+ \int_0^t ds \left\{ \sup_{\substack{f \geq 0 \\ \int f d\nu_\alpha^N = 1}} \left\{ \int Z_{N,\varepsilon}^1(\pm G_s, \eta) f(\eta) d\nu_\alpha^N(\eta) + \frac{N^{2-d}}{A} \left\langle \sqrt{f}, \mathbf{L}_\gamma^0 \sqrt{f} \right\rangle_\alpha \right\} \right\}. \end{aligned}$$

It has been proven in [KLO], [KL] that for any positive  $A$  and any smooth function  $G \in \mathcal{C}^{1,1}([0, T] \times \mathbf{T}^d)$  and any  $\alpha$

$$\lim_{n \rightarrow \infty} \limsup_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} \int_0^t ds \left\{ \sup_{\substack{f \geq 0 \\ \int f d\nu_\alpha^N = 1}} \left\{ \int Z_{N,\varepsilon}^1(G_s, \eta) f(\eta) d\nu_\alpha^N(\eta) + AN^{2-d} \left\langle \sqrt{f}, \mathbb{L}_\gamma^0 \sqrt{f} \right\rangle_\alpha \right\} \right\} = 0.$$

That concludes the proof of Lemma 6.5.  $\square$

*Proof of Lemma 6.6.* Fix  $0 \leq k \leq d$ . To simplify the notation, for  $0 \leq m \leq d$  denote by  $\psi^1$  and  $\psi^{2,m}$  the cylinder functions  $\Psi^1 = r_{0,e_1}$  and

$$\Psi_k^{2,m} = \left\{ \sum_{j \in \mathbf{T}^d} (1/2) (\nabla_{j,j+e_m} F^{n,k} - \nabla_{j+e_m,j} F^{n,k}) \right\}.$$

For a cylinder function  $\Psi$  and a positive density  $0 \leq \rho \leq 2$ , we shall denote by  $\tilde{\Psi}(\rho)$  its expectation with respect to the measure  $\nu_\rho^N$ .

Observe that for each  $0 \leq m \leq d$  and all density  $0 \leq \rho \leq 2$ ,  $\tilde{\Psi}_k^{2,m}(\rho)$  can be rewritten as

$$\begin{aligned} \tilde{\Psi}_k^{2,m}(\rho) &= - \langle 2W_{0,e_m}, F^{n,k} \rangle_{\rho,0} \\ &= \lll 2W_{0,e_m}, \mathbb{L}^0 F^{n,k} \ggg_\rho. \end{aligned}$$

In particular, from Lemma 6.8 and using one and two blocks estimates, we obtain

$$\left| \int_0^T Z_{N,\varepsilon}^{F^{k,n},2}(G_s, \eta_s) ds \right| = \left| \int_0^T \tilde{Z}_{N,\varepsilon}^{F^{k,n},2}(G_s, \eta_s) ds \right| + r_{N,\varepsilon},$$

where  $r_{N,\varepsilon}$  is a random variable such that its expectation with respect to  $P_{\mu^N}$  converges to 0 as  $N \uparrow \infty$  and  $\varepsilon \downarrow 0$  and

$$\begin{aligned} \tilde{Z}_{N,\varepsilon}^{F^{k,n},2}(G, \eta) &= \beta N^{-d} \sum_{i \in \mathbf{T}^d} G(i/N) \left\{ \sum_{m=1}^d \left( \partial_m J * \pi^N(\eta)(i/N) \right) \left\{ \tilde{\Psi}^1(\eta^{N\varepsilon}(i)) \delta_{k,m} \right. \right. \\ &\quad \left. \left. - 2(\chi D_{k,m})(\eta^{N\varepsilon}(i)) - \tilde{\Psi}_k^{2,m}(\eta^{N\varepsilon}(i)) \right\} \right\}. \end{aligned}$$

To conclude the proof we just have to apply Lemma 6.7 below.  $\square$

**Lemma 6.7** *There exists positive constant  $C_0$  such that for each  $0 \leq m \leq d$*

$$\sup_{0 \leq \rho \leq 2} \left| \langle r_{0,e_1} \rangle_\rho \delta_{k,m} - 2\chi(\rho) D_{k,m}(\rho) - 2 \lll W_{0,e_m}, \mathbb{L}^0 F^{k,n} \ggg_\rho \right| \leq \frac{C_0}{\sqrt{n}}.$$

*Proof.* Using (6.10), it is easy to show that

$$\begin{aligned} & \left| \langle r_{0,e_1} \rangle_\rho \delta_{k,m} - 2\chi(\rho)D_{k,m}(\rho) - 2\ll W_{0,e_m}, \mathbb{L}^0 F^{k,n} \gg_\rho \right| \\ &= \left| \ll 2W_{0,m}, W_{0,e_k} + \sum_{r=1}^d D_{k,r}(\rho) [\eta(e_r) - \eta(0)] - \mathbb{L}^0 F^{k,n} \gg_\rho \right|. \end{aligned}$$

By Schwartz inequality the right hand side of the last expression is bounded by

$$\sqrt{\ll 2W_{0,m} \gg_\rho} \times \left( \ll W_{0,e_k} + \sum_{r=1}^d D_{k,r}(\rho) [\eta(e_r) - \eta(0)] - \mathbb{L}^0 F^{k,n} \gg_\rho \right)^{\frac{1}{2}}$$

which is bounded by  $C_0/\sqrt{n}$  for some positive constant  $C_0$ .  $\square$

We now prove some estimates on the Dirichlet forms for the process at  $\beta \neq 0$  needed in the proof of Theorem 6.4 and in the proof of one and two blocs estimates.

**Lemma 6.8** *There exists positive constant  $C_1$  such that for every probability density  $f$  with respect to  $\nu_\alpha^N$*

$$N^2 \left\langle \sqrt{f}, \mathbb{L}_\gamma^\beta \sqrt{f} \right\rangle_{\nu_\alpha^N} \leq \frac{1}{2} N^2 \left\langle \sqrt{f}, \mathbb{L}_\gamma^0 \sqrt{f} \right\rangle_{\nu_\alpha^N} + C_1 N^d.$$

*Proof.* Fix a probability density  $f$ . A simple computation shows that

$$\begin{aligned} N^2 \left\langle \sqrt{f}, \mathbb{L}_\gamma^\beta \sqrt{f} \right\rangle_{\nu_\alpha^N} &= N^2 \left\langle \sqrt{f}, \mathbb{L}_\gamma^0 \sqrt{f} \right\rangle_{\nu_\alpha^N} \\ &+ \frac{1}{2} \sum_{\substack{i,j \in \mathbf{T}^d \\ |i-j|=1}} \int r_{i,j} \left[ 1 - \Phi \left\{ \beta (\nabla^{i,j} H_N) (\eta) \right\} \right] \sqrt{f} \left[ \nabla_{i,j} \sqrt{f} \right] (\eta) d\nu_\alpha^N. \end{aligned} \quad (6.14)$$

Recall that, by Lemma 6.9 below,

$$\left| 1 - \Phi \left\{ \beta (\nabla^{i,j} H_N) (\eta) \right\} \right| \leq C_2 N^{-1}$$

for some positive constant  $C_2$ . To conclude the proof of the lemma it remains to bound the second term of the right hand side of (6.14) by

$$-(N^2/2) \left\langle \sqrt{f}, \mathbb{L}_\gamma^0 \sqrt{f} \right\rangle_{\nu_\alpha^N} + C_2 N^d$$

using the inequalities  $2xy \leq ax^2 + a^{-1}y^2$  for all  $a > 0$  and the fact that  $f$  is a density with respect to  $\nu_\alpha^N$ .  $\square$

**Lemma 6.9** For all  $i \in \mathbf{T}^d$ , unit vector  $e \in \mathbf{T}^d$  and  $\eta \in \mathbb{X}_N$

$$C_\gamma^\beta(i, i + e; \eta) = 1 + \left\{ \beta N^{-d-1} \sum_{j \in \mathbf{T}^d} (e \cdot \nabla J) ((i - j)/N) \eta(j) \right\} + O(N^2).$$

*Acknowledgments.* We would like to thank R. Esposito and E. Presutti for useful discussions. R. M. is indebted to G.B. Giacomin for sending his preliminary notes on a related lattice gas model. R. M. would like to thank J. L. Lebowitz for bringing to her attention the Blume-Capel model. M.M. would like to thank the very kind hospitality at the University of Roma Tor Vergata and R. M. the hospitality of the University of Rouen.

## Appendix.

**Proof of (5.1)** We have

$$\begin{aligned} \sum_{k=1}^d |\partial_k h_{t+\varepsilon-s}(x)| &= (2\pi(t+\varepsilon-s))^{-d/2} \exp \left\{ -\frac{1}{8(t+\varepsilon-s)} \|x\|^2 \right\} \times \frac{1}{\sqrt{t+\varepsilon-s}} \times \\ &\quad \times \frac{\sum_{k=1}^d |x_k|}{\sqrt{t+\varepsilon-s}} \exp \left\{ -\frac{1}{8(t+\varepsilon-s)} \|x\|^2 \right\}, \end{aligned}$$

where for  $x \in \mathbb{R}^d$ ,  $\|x\| = \sqrt{\sum_{k=1}^d (x_k)^2}$ . Since  $\sum_{k=1}^d |x_k| \leq d\|x\|$  and  $\alpha \exp(-\alpha^2) \leq A_1$  for all  $\alpha \in \mathbb{R}$  and for some positive constant  $A_1$ . We have that

$$\sum_{k=1}^d |\partial_k h_{t+\varepsilon-s}(x)| \leq A_2 h_{2(t+\varepsilon-s)}(x) \frac{1}{\sqrt{t+\varepsilon-s}}$$

for some positive constant  $A_2$  that depends on  $d$ . On the other hand, we have

$$\sum_{k=1}^d \partial_k \mathcal{H}_{t,\varepsilon}^f = f * \left( \sum_{k=1}^d \partial_k h_{t+\varepsilon-s} \right)$$

and therefore

$$\begin{aligned}
& \sum_{k=1}^d \int_0^t \left| \langle \partial_k \mathcal{H}_{t,\varepsilon}^f(s, \cdot) \rangle \right| ds \\
& \leq \int_0^t ds \left\{ \int_{\mathbb{R}^d} dx \left[ \int_{\mathbb{R}^d} |f(y)| \sum_{k=1}^d |\partial_k h_{t+\varepsilon-s}(y-x)| dy \right] \right\} \\
& \leq A_2 \int_0^t ds \left\{ \int_{\mathbb{R}^d} dy |f(y)| \left[ \int_{\mathbb{R}^d} h_{2(t+\varepsilon-s)}(y-x) dx \right] \times \frac{1}{\sqrt{t+\varepsilon-s}} \right\} \\
& \leq A_2 \|f\|_1 \int_0^t \frac{1}{\sqrt{t+\varepsilon-s}} ds \\
& = A_2 \|f\|_1 \left\{ \sqrt{t+\varepsilon} - \sqrt{\varepsilon} \right\}.
\end{aligned}$$

□

**Proof of (5.3):** For all  $\tau \in [0, t]$ ,  $W(\tau, \cdot) \in L^\infty(\mathbb{R}^d)$  and  $\|W(\tau, \cdot)\|_\infty \leq 2C_3\sqrt{t}R(t)$ .

First, by (5.2), for any open set  $U$  of  $\mathbb{R}^d$  with finite Lebesgue measure  $\lambda(U)$ , we have for all  $0 \leq \tau \leq t$ ,

$$\int_U W(\tau, x) dx \leq 2C_3\sqrt{t}R(t)\lambda(U).$$

Fix  $0 < \delta < 1$ . For any open set  $U$  of  $\mathbb{R}^d$  with finite Lebesgue measure and for  $0 \leq \tau \leq t$  let

$$B_{\delta,\tau}^U = \left\{ x \in U : W(\tau, x) > 2C_3\sqrt{t}R(t)(1+\delta) \right\}.$$

Suppose that  $\lambda(B_{\delta,\tau}^U) > 0$ , there exists an open set  $V$ , such that,  $B_{\delta,\tau}^U \subset V$  and  $\lambda(V \setminus B_{\delta,\tau}^U) \leq \lambda(V)\frac{\delta}{2}$  and we have

$$\begin{aligned}
\lambda(V)(2C_3\sqrt{t}R(t)) & < \lambda(V)(2C_3\sqrt{t}R(t))(1+\delta)(1-\delta/2) \\
& = (2C_3\sqrt{t}R(t))(1+\delta)(\lambda(V) - \lambda(V)\delta/2) \\
& \leq (2C_3\sqrt{t}R(t))(1+\delta)(\lambda(V) - \lambda(V \setminus B_{\delta,\tau}^U)) \\
& = (2C_3\sqrt{t}R(t))(1+\delta)\lambda(B_{\delta,\tau}^U) \\
& < \int_{B_{\delta,\tau}^U} W(\tau, x) dx \\
& \leq \int_V W(\tau, x) dx \\
& \leq (2C_3\sqrt{t}R(t))\lambda(V).
\end{aligned}$$

which leads to a contradiction.



By the arbitrariness of  $0 < \delta < 1$  we obtain that if  $U$  is any open set of  $\mathbb{R}^d$  with  $\lambda(U) < \infty$ ,

$$\lambda\left(\left\{x \in U : W(\tau, x) > 2C_3\sqrt{t}R(t)\right\}\right) = 0.$$

This implies

$$W(\tau, x) \leq 2C_3\sqrt{t}R(t) \quad \text{a.e. in } \mathbb{R}^d.$$

□

## References

- [B] Blume, M, "Theory of the First Order Magnetic Phase Change in  $\text{UO}_2$ " *Phys. Rev.* **141**, 517, (1996)
- [C] Capel, H.W., *Physica* **32**, 96 (1996).
- [BEG] Blume, M., Emery, V. J. and Griffiths R. B. "Ising Model for the  $\lambda$  Transition and Phase Separation in  $\text{He}^3$ - $\text{He}^4$  Mixtures". *Phys. Rev. A*, **4**, 1071-1077, (1971).
- [BS] Bricmont, J. and Slawny, J., "Phase transition in Systems with a Finite Number of dominant Ground States", *Journ. Stat. Phys.*, **54**, 89, (1989).
- [CCO] Carlen, E. A., Carvalho, M.C. and Orlandi, E. "Algebraic rate of decay for the excess free energy and stability of fronts for a non-local phase kinetics equation with a conservation law", I. to appear on *J.Stat.Phys.* **95** (1999)  
"Algebraic rate of decay for the excess free energy and stability of fronts for a non-local phase kinetics equation with a conservation law, II". to appear on *Comm. Par. Diff. Eq.* (1999).
- [DOPT] De Masi, A., Orlandi, E., Triolo, L. and Presutti, E. "Glauber evolution with Kac potentials I. Mesoscopic and macroscopic limits, interface dynamics", *Nonlinearity* **7**, 287–301, (1996).  
"Glauber evolution with Kac potentials II. Fluctuation". *Nonlinearity* **9**, 27–51, (1996).  
"Glauber evolution with Kac potentials. III. Spinodal decomposition". *Nonlinearity* **9**, 53–114, (1996).
- [DP] De Masi, A. and Presutti, E. *Mathematical Methods for Hydrodynamic limits*, *Lecture Notes in Mathematics*, **1501**, Springer-Verlag, Berlin, (1991).
- [F] Fritz, J. "On the diffusive nature of the entropy flow in infinite systems: remarks to a paper by Guo-Papanicolau-Varadhan". *Commun. Math. Phys.* **133**, 331-352. (1990)
- [GL] Giacomini, G. and Lebowitz, J. L. "Phase segregation dynamics in particle systems with long range interaction I. Macroscopic limits". *J. Stat. Phys.* **87**, 37-61. (1997).  
"Phase segregation dynamics in particle systems with long range interaction II. Interface motion", *SIAM J. Appl. Math.* **58**, 1707–1729 (1998).
- [GLM] Giacomini, G., Lebowitz, J. L. and Marra, R. in preparation.

- [GLP] Giacomini, G., Lebowitz, J. L. and Presutti, E. "Deterministic and Stochastic Hydrodynamic Equations Arising From Simple Microscopic Model Systems". Preprint (1998).
- [GPV] Guo, M.Z., Papanicolaou, G.C. and Varadhan, S.R.S. "Nonlinear diffusion limit for a system with nearest neighbor interactions", *Comm. Math. Phys.*, **118**, 31-59. (1988).
- [KUE] Kac, M., Uhlenbeck, G.E. and Hemmer, P.C. "On the van der Waals theory of the vapor-liquid equilibrium, I. Discussion of a one-dimensional model", *J. Math. Phys.* **4**, 216–228, (1963).
- [KL] Kipnis, C. and Landim, C. *Scaling limits of interacting particle systems*, Springer-Verlag, (1999).
- [KLO1] C. Kipnis, C. Landim and S. Olla, "Hydrodynamical limit for a nongradient model: the generalized exclusion processes", *Comm. Pure Appl. Math.*, **47**, 1994, pp. 1475–1545.
- [KOV] Kipnis, C., Olla, S. and Varadhan, S.R.S. Hydrodynamics and large deviations for simple exclusion processes, *Comm. Pure Appl. Math.* **42**, 115-137. (1989).
- [Li] Liggett, T. M. Interacting particle systems. **276** Grundlehren der mathematischen Wissenschaften. Springer-Verlag. Berlin. (1985).
- [LM] Landim, C. and Mourragui, M. "Hydrodynamic limit of mean zero asymmetric zero range processes in infinite volume". *Ann.Inst.Henri.Poincaré*. Vol 33, N. **1**, p.361-385, (1997).
- [LMS] C. Landim, M. Mourragui and S. Sellami, "Hydrodynamical limit for a nongradient interacting particle system with stochastic reservoirs". Preprint (1998)
- [LP] Lebowitz, J. and Penrose, O. "Rigorous treatment of the Van der Waals Maxwell theory of the liquid vapour transition", *J. Math. Phys.* **7**, 98–113 (1963)
- [Q] Quastel, J. "Diffusion of a color in the simple exclusion process", *Comm. Pure Appl. Math.* **XLV**, 623–679 (1992).
- [Sp] Spohn, H. Large scale dynamics of interacting particles, Springer, Berlin, (1991).
- [V] S.R.S. Varadhan, "Nonlinear diffusion limit for a system with nearest neighbor interactions II", in *Asymptotic Problems in Probability Theory: Stochastic Models and Diffusion on Fractals*, edited by K. Elworthy and N. Ikeda, Pitman Research Notes in Mathematics 283, Wiley, (1994).
- [VY] S. R. S. Varadhan and H.T. Yau "Diffusive Limit of Lattice Gas with Mixing Conditions. *Asian J. Math.* **1**, 623–678 (1997).
- [Y] Yau, H. T. "Metastability of Ginzburg-Landau model with a conservation law", *J. Stat. Phys.* **74** 705-742. (1994).